

2013 Texas Childcare Immunization Assessment Coverage Level Results

The Texas Childcare Immunization Assessment survey is conducted triennially to assess vaccination coverage levels of children attending a childcare facility and to monitor compliance with new and existing vaccine requirements at licensed child care centers (LCCC) and registered child care homes (RCCH) in Texas. A random sample of childcare facilities is selected for the survey. Local health departments and Texas Department of State Health Services staff review a random sample of immunization records from each selected childcare facility. Statewide and regional vaccination coverage levels are available for children ages 19 to 59 months of age attending childcare facilities in Texas. The percent of children aged 19 to 59 months that attend childcare and have an exemption to one or more vaccines was 0.42%.

Vaccination Coverage Levels among Childcare Attendees Aged 19 – 59 Months, Texas and Health Service Regions (HSR), 2013

Area	DTaP‡	Polio§	MMR	Var††	HepB*	Hib¶	PCV§§	HepA**	4:3:1:3:3:1:4^
Texas	90.5	97.0	95.2	94.9	93.4	97.1	88.2	73.5	78.9
HSR 1	93.2	97.9	96.5	96.2	94.0	97.8	89.2	79.0	79.7
HSR 2/3	88.0	97.5	93.4	93.1	91.0	97.4	86.7	71.3	75.1
HSR 4/5N	92.5	97.6	97.0	96.8	95.7	97.4	89.8	74.5	84.0
HSR 6/5S	92.5	97.0	95.7	95.8	94.8	97.2	89.8	71.9	81.7
HSR 7	87.2	95.1	93.0	92.6	90.0	94.9	83.8	69.4	73.2
HSR 8	91.1	96.5	94.0	94.2	95.1	97.9	89.6	72.5	79.1
HSR 9/10	89.6	97.1	95.6	95.2	92.2	97.0	87.8	76.9	78.6
HSR 11	90.3	96.8	95.9	95.5	93.0	97.5	89.3	78.4	78.5

‡ 4 or more doses of any diphtheria and tetanus toxoids and pertussis vaccines including diphtheria and tetanus toxoids, and any acellular pertussis vaccine (DTaP/DTP/DT).

§ 3 or more doses of any poliovirus vaccine.

|| 1 or more doses of measles-mumps-rubella vaccine.

¶ 3 or more doses of Haemophilus influenzae type b (Hib) vaccine.

* 3 or more doses of hepatitis B vaccine.

†† 1 or more doses of varicella at or after child's first birthday, unadjusted for history of varicella illness.

§§ ≥4 doses of pneumococcal conjugate vaccine (PCV).

** 2 or more doses of hepatitis A vaccine.

^ ≥4 doses of diphtheria and tetanus toxoids and acellular pertussis (DTaP) vaccine, ≥3 doses of poliovirus (Polio) vaccine, ≥1 dose of any measles-mumps-rubella (MMR) vaccine, ≥3 doses of Haemophilus influenzae (Hib) vaccine of any type, ≥3 doses of hepatitis B (HepB) vaccine, ≥1 dose of varicella (Var) vaccine, and ≥4 doses of pneumococcal conjugate vaccine (PCV).