

NPI NorthSTAR Dx LAYOUT 2/1/07

Field	Len	Begin	End	Comment/Decode
CMP	3			Header Information to be stripped out
SEMI_COL1	1			Header Information to be stripped out
RPT_NM_WITH_YYJJJ	12			Header Information to be stripped out
SEMI_COL2	1		17	Header Information to be stripped out
REC_TYPE	12	1	12	Value = "DIAG"
EFF_DT	8	13	20	Decision Date which is the date the client's diagnosis was determined.
ID	10	21	30	Identifies a client's CARE ID.
COMP	3	31	33	Component code.
NS_COMP	3	34	36	NorthSTAR component code.
NS_ID	10	37	46	NorthSTAR ID, if fewer than 10 digits, it will be padded with leading zeros to a length of 10.
MHPRIPOP	1	47	47	Indicates whether the diagnosis places the person in the MH priority population. Acceptable values are Y or N.
MHPRIPOP_REAS	6	48	53	Indicates a reason a person is not in the MH priority population. See CARE reference manual for decodes.
REAS_ACT	1	54	54	1=Admission, 2=Reevaluation, 3 = Death, 4 = Discharge.
AXIS_IND	1	55	55	Indicates which axis (I or II) a client's principal diagnosis is determined.
FORM_TM	4	56	59	Time recorded on the CARE -DG-1 form. HHMM military time format.
DSM11	5	60	64	DSM code for psychiatric syndromes diagnoses. Level 1 indicates the most significant diagnosis, level 6 the least significant. See CARE reference manual for decodes.
DSM12	5	65	69	DSM code for psychiatric syndromes diagnoses. Level 1 indicates the most significant diagnosis, level 6 the least significant. See CARE reference manual for decodes.
DSM13	5	70	74	DSM code for psychiatric syndromes diagnoses. Level 1 indicates the most significant diagnosis, level 6 the least significant. See CARE reference manual for decodes.
DSM14	5	75	79	DSM code for psychiatric syndromes diagnoses. Level 1 indicates the most significant diagnosis, level 6 the least significant. See CARE reference manual for decodes.
DSM15	5	80	84	DSM code for psychiatric syndromes diagnoses. Level 1 indicates the most significant diagnosis, level 6 the least significant. See CARE reference manual for decodes.
DSM16	5	85	89	DSM code for psychiatric syndromes diagnoses. Level 1 indicates the most significant diagnosis, level 6 the least significant. See CARE reference manual for decodes.

DSM21	5	90	94	DSM code for personality, specific developmental disorders and mental retardation. Level 1 indicates the most significant diagnosis, level 4 the least significant.
DSM22	5	95	99	DSM code for personality, specific developmental disorders and mental retardation. Level 1 indicates the most significant diagnosis, level 4 the least significant. See CARE reference manual for decodes.
DSM23	5	100	104	DSM code for personality, specific developmental disorders and mental retardation. Level 1 indicates the most significant diagnosis, level 4 the least significant.
DSM24	5	105	109	DSM code for personality, specific developmental disorders and mental retardation. Level 1 indicates the most significant diagnosis, level 4 the least significant. See CARE reference manual for decodes.
DSM31	5	110	114	Axis 3 level 1. ICD-9 code for physical disorders. Level 1 indicates the most significant diagnosis, level 6 the least significant. See 'ICDCODES' decode table conditionally required: if principal diagnosis = MH or reason for action = 3(death),
				then Axis III Level I ICD code is required. If no Axis II DX, then use the word 'NONE' to indicate that there is no diagnosis, but that the diagnosis was reviewed at the time of update. Data entry is a required field at update.
DSM32	5	115	119	ICD-9 code for physical disorders. Level 1 indicates the most significant diagnosis, level 6 the least significant. See CARE reference manual for decodes.
DSM33	5	120	124	ICD-9 code for physical disorders. Level 1 indicates the most significant diagnosis, level 6 the least significant. See CARE reference manual for decodes.
DSM34	5	125	129	ICD-9 code for physical disorders. Level 1 indicates the most significant diagnosis, level 6 the least significant. See CARE reference manual for decodes.
DSM35	5	130	134	ICD-9 code for physical disorders. Level 1 indicates the most significant diagnosis, level 6 the least significant. See CARE reference manual for decodes.
DSM36	5	135	139	ICD-9 code for physical disorders. Level 1 indicates the most significant diagnosis, level 6 the least significant. See CARE reference manual for decodes.
AXIS3_DT	8	140	147	Date of the physician's examination in which the Axis 3 diagnosis was determined. Format = YYYYMMDD. Must be < = decision date. No date required if only Axis III entry is NONE.
AXIS4	9	148	156	Code for psychosocial stressors. There are up to 9 occurrences (stressors). See CARE reference manual for decodes.

AXIS5	2	157	158	Indicates a clinician's overall judgment of a person's psychological, social, and occupational functioning using the Global Assessment of Functioning Scale (GAF Scale) that assess mental illness. See CARE reference manual for decodes.
AXIS5_PREV	2	159	160	The previous AXIS 5 code.
ABL_CUR	1	161	161	Identifies a person's potential adaptive behavior level.
ABL_POT	1	162	162	Identifies a client's current adaptive behavior level. See CARE reference manual for decodes.
AAMD_PRIM	3	163	165	Code to indicate a person's primary AAMD disorder, if one exists. For MR persons only. See CARE reference manual for decodes.
AAMD_SEC	3	166	168	Code to indicate a client's secondary AAMD disorder, if one exists. For MR persons only.
AAMD_TERT	3	169	171	Code to indicate a person's tertiary AAMD disorder, if one exists. For MR persons only. See CARE reference manual for decodes.
GEN_COMP	2	172	173	Code to indicate whether a person has a genetic defect. See CARE reference manual for decodes.
CRAN_ANOM	2	174	175	Code to indicate whether a person has a cranial anomaly. See CARE reference manual for decodes.
SENS_IMPAIR	2	176	177	Code to indicate whether a person has a sensory impairment. See CARE reference manual for decodes.
PERCEPT	2	178	179	Code to indicate whether a person has a perception disorder. See CARE reference manual for decodes.
CONVULS_DIS	2	180	181	Code to indicate whether a person has a convulsive disorder. See CARE reference manual for decodes.
PSY_IMPAIR	2	182	183	Code to indicate whether a person has a psychiatric impairment. See CARE reference manual for decodes.
MOTOR_DYS	2	184	185	Code to indicate a person's motor dysfunction type. See CARE reference manual for decodes.
MOTOR_LOC	1	186	186	Code to indicate a person's motor dysfunction location. See CARE reference manual for decodes.
MOTOR_SEV	1	187	187	Code to indicate a person's motor dysfunction severity. See CARE reference manual for decodes.
AAMD_DT	8	188	195	Date of the physician's examination in which the AAMD diagnoses were determined. Format = YYYYMMDD.
DSM_VERS	1	196	196	Version of the DSM codes used for diagnosis.
ICD_VERS	1	197	197	Version of the ICD codes used for diagnosis.
AADM_VERS	2	198	199	Version of the AAMD codes used for diagnosis.
IQ	3	200	202	A person's IQ score.
IQ_TEST_DT	8	203	210	Date of the IQ test.
IQ_TEST	2	211	212	Type of IQ test. See CARE reference manual for decodes.
SQ	3	213	215	A person's SQ score.
SQ_TEST_DT	8	216	223	Date of the SQ test.
SQ_TEST	2	224	225	Type of SQ test. See CARE reference manual for decodes.
ADD_DT	8	226	233	Mainframe housekeeping fields

LASTUP_DT	8	234	241	Mainframe housekeeping fields
LASTUP_ID	8	242	249	Mainframe housekeeping fields
LASTUP_TM	4	250	253	Mainframe housekeeping fields
LOC_PROV_ID	9	254	262	Location/provider ID. The current input for this field is 6 characters. Nine field length has been allowed for expansion. The field will be read from left to right.
NPI	10	263	272	Provider's NPI