

## COMFORT MEASURES

Return for next immunization visit \_\_\_\_\_

### INJECTION SITE

**L R DTaP, DT, Td or Tdap** – Diphtheria, Tetanus, Pertussis (Whooping Cough)  
\_\_\_\_\_ During the first 24 hours, a slight fever may develop (101° under the arm, 102° rectal), irritability, or local redness and/or tenderness at the site of the injection, prior to or immediately after immunization. Give extra fluids. Bathe in lukewarm water as often as needed. Symptoms may continue for up to 3 days.

**L R IPV** – Inactivated Polio  
\_\_\_\_\_ Tenderness at injection site

**L R MMR, Varicella, Zoster** – Measles, Mumps, Rubella, Varicella (Chicken Pox), Zoster (Shingles)  
\_\_\_\_\_ **Measles** – Some develop a rash and/or a slight fever 1-2 weeks following immunization. The rash is not contagious. Treat like any mild fever (extra fluids)  
**Mumps** - On rare occasions, the salivary glands may become swollen within 4 weeks following immunization. No specific treatment.  
**Rubella** – Some may develop a rash or mild swelling of the neck glands 1-2 weeks after the vaccine. Some may also experience joint swelling or aching 1-3 weeks after receiving the vaccine. If joint swelling or aching is severe, see your doctor.  
**Varicella** -Tenderness at site of injection, slight fever (treat with extra fluids), mild rash may develop up to a month after receiving vaccination. This rash is contagious and others in the family may become infected.  
**Zoster** - Mild headache, redness, soreness, swelling or itching at site.

**L R Hib** – Haemophilus Influenzae Type B  
\_\_\_\_\_ Tenderness at the site of injection and/or slight fever. Treat like any mild fever (extra fluids).

**L R HPV** – Human Papillomavirus  
\_\_\_\_\_ Tenderness at the site of injection and/or slight fever. Treat like any mild fever (extra fluids).

**L R Hep A or Hep B** – Hepatitis A, Hepatitis B  
\_\_\_\_\_ Tenderness at the site of injection and/or slight fever. Treat like any mild fever (extra fluids).

**L R PCV13, PPV23** – Pneumococcal  
\_\_\_\_\_ During the first 24-48 hours, tenderness or redness at the site of injection, slight fever (extra fluids)

**L R MCV4** – Meningococcal Conjugate  
\_\_\_\_\_ During the first 24-48 hours, tenderness or redness at the site of injection.

**L R Flu** – influenza  
\_\_\_\_\_ Redness, swelling, tenderness at site of injection, aches, fever (extra fluids).

**ORAL Rotavirus**  
\_\_\_\_\_ Within 7 days of ingestion may experience mild, temporary diarrhea or vomiting.

**If the receiver of the vaccine becomes ill and visits a doctor, hospital or clinic during the four weeks after receiving immunization(s), please report it to our office: \_\_\_\_\_ OR contact the Immunization Division at the Texas Department of State Health Services at 2408 South 37<sup>th</sup> Street, Temple, Texas 76504, (254) 778-6744.**