

Fallopian Tube**C57.0**

C57.0 Fallopian tube

Note: Laterality must be coded for this site.**FallopianTube****CS Tumor Size****See Standard Table****FallopianTube****CS Extension****Note 1:** Positive regional lymph nodes (FIGO Stage IIIC) are coded in the CS Lymph Nodes field.**Note 2:** Codes 130 and 660: Since "malignant ascites or malignant peritoneal washings" was not specifically categorized in the 1977 Summary Staging Guide, it is unclear to which stage previous cases may have been coded.**Note 3:** Liver capsule metastases are coded to 750-780 in the Extension field; liver parenchymal metastases are coded in the Mets at DX field.**Note 4:** Both extension to and discontinuous metastasis to any of the following PELVIC organs is considered FIGO Stage II and coded in the range 350-660: adnexae, NOS; bladder, bladder serosa; broad ligament (mesovarium); cul-de-sac; fallopian tubes; parametrium; pelvic peritoneum; pelvic wall; rectum; sigmoid colon; sigmoid mesentery; ureter; uterus; uterine serosa.**Note 5:** Both extension to and discontinuous metastasis to any of the following ABDOMINAL organs is considered FIGO Stage III and coded in the range 690-780: abdominal mesentery; diaphragm; gallbladder; infracolic omentum; kidneys; large intestine except rectum and sigmoid colon; liver (peritoneal surface); omentum; pancreas; pericolic gutter; peritoneum, NOS; small intestine; spleen; stomach; ureters.**Note 6:** From the AJCC Manual 6th Edition (page 285): "It may be preferable to classify a patient as TX (primary tumor cannot be assessed) if inadequate staging biopsies and/or a lack of peritoneal cytology make it inaccurate to classify the patient with confidence as early stage (Stage T3a/IIIA has not been excluded by adequate staging biopsies)."

Code	Description	TNM 7 Map	TNM 6 Map	SS77 Map	SS2000 Map
000	In situ: noninvasive, intraepithelial Limited to tubal mucosa FIGO Stage 0	Tis	Tis	IS	IS
100	Confined to fallopian tube, NOS FIGO Stage I	T1NOS	T1NOS	L	L
110	Confined to one fallopian tube WITHOUT penetrating serosal surface; no ascites	T1a	T1a	L	L

Code	Description	TNM 7 Map	TNM 6 Map	SS77 Map	SS2000 Map
110 cont'd	FIGO Stage IA	T1a	T1a	T1a	L
120	Confined to both fallopian tubes WITHOUT penetrating serosal surface; no ascites FIGO Stage IB	T1b	T1b	L	L
130	Extension onto or through tubal serosa Malignant ascites Malignant peritoneal washings FIGO Stage IC	T1c	T1c	L	L
300	Localized, NOS FIGO Stage I	T1NOS	T1NOS	L	L
350	Pelvic extension, NOS with no malignant cells in ascites or peritoneal washings FIGO Stage II	T2NOS	T2NOS	RE	RE
400	Extension or metastasis to (but no malignant cells in ascites or peritoneal washings): Corpus uteri Ovary, ipsilateral Uterus, NOS FIGO Stage IIA	T2a	T2a	RE	RE
450	Ovary, contralateral (but no malignant cells in ascites or peritoneal washings)	T2a	T2a	D	RE
460	FIGO Stage IIA	T2a	T2a	RE	RE
500	Extension or metastasis to (but no malignant cells in ascites or peritoneal washings): Broad ligament, ipsilateral Mesosalpinx, ipsilateral Adjacent peritoneum	T2b	T2b	RE	RE

Code	Description	TNM 7 Map	TNM 6 Map	SS77 Map	SS2000 Map
500 cont'd	FIGO Stage IIB	T2b	T2b	RE	RE
600	OBSOLETE DATA RETAINED V0200 Ovary, contralateral (but no malignant cells in ascites or peritoneal washings) FIGO Stage IIA	ERROR	T2a	D	RE
610	OBSOLETE DATA RETAINED V0200 (600) + (500)	ERROR	T2b	D	RE
620	(500) + (450)	T2b	T2b	D	RE
650	Extension or metastasis to (but no malignant cells in ascites or peritoneal washings): Broad ligament, contralateral Cul de sac (rectouterine pouch) Mesosalpinx, contralateral Rectosigmoid Sigmoid FIGO IIB	T2b	T2b	D	RE
660	Pelvic extension (codes 350-650) WITH malignant cells in ascites or peritoneal washings FIGO Stage IIC	T2c	T2c	D	D
680	Peritoneal implants or metastasis(size of metastases not stated; unknown if microscopic or macroscopic) Omentum Small intestine FIGO Stage III	T3NOS	T3NOS	D	RE
690	Microscopic peritoneal implants or metastasis: Omentum Small intestine	T3a	T3a	D	RE

Code	Description	TNM 7 Map	TNM 6 Map	SS77 Map	SS2000 Map
690 cont'd	FIGO Stage IIIA	T3a	T3a	D	RE
700	OBSOLETE DATA CONVERTED V0102 See code 680 Extension or metastasis to: Omentum FIGO Stage IIB	ERROR	ERROR	D	RE
710	OBSOLETE DATA CONVERTED V0102 See code 660 Pelvic extension (codes 350-700) with malignant cells in ascites or peritoneal washings FIGO Stage IIC	ERROR	ERROR	D	D
720	Macroscopic peritoneal implants or metastasis less than or equal to 2cm Omentum Small intestine FIGO Stage IIIB	T3b	T3b	D	RE
730	Macroscopic peritoneal implants or metastasis greater than 2cm Omentum Small intestine FIGO Stage IIIC	T3c	T3c	D	RE
750	Peritoneal implants outside the pelvis, NOS (size of metastases not stated; unknown if microscopic or macroscopic), except code 680 (See Note 5)	T3NOS	T3NOS	D	D
760	Microscopic peritoneal metastasis outside the pelvis, except code 690 (See Note 5) FIGO Stage IIIA	T3a	T3a	D	D

Code	Description	TNM 7 Map	TNM 6 Map	SS77 Map	SS2000 Map
770	Macroscopic peritoneal metastasis less than or equal to 2 cm outside the pelvis, except code 720 (See Note 5) FIGO Stage IIIB	T3b	T3b	D	D
780	Peritoneal metastases greater than 2 cm, except code 730 (See Note 5) FIGO Stage IIIC	T3c	T3c	D	D
800	Further contiguous extension FIGO Stage III [NOS]	T3NOS	T3NOS	D	D
950	No evidence of primary tumor	T0	T0	U	U
999	Unknown extension Primary tumor cannot be assessed (See Note 6) Not documented in patient record	TX	TX	U	U

FallopianTube**CS Tumor Size/Ext Eval**

See Standard Table

FallopianTube**CS Lymph Nodes**

Note 1: Code only regional nodes and nodes, NOS, in this field. Distant nodes are coded in the field Mets at DX.

Note 2: If the clinician says "adnexa palpated" but doesn't mention lymph nodes, assume lymph nodes are not involved.

Note 3: If either exploratory/definitive surgery is done with no mention of lymph nodes, assume nodes are negative.

Note 4: Regional nodes include bilateral and contralateral involvement of named nodes.

Note 5: Positive presacral and lateral sacral (laterosacral) lymph nodes are coded under CS Mets at DX.

Note 6: Determination of the N category is based on this field and CS Mets at DX.

Code	Description	TNM 7 Map	TNM 6 Map	SS77 Map	SS2000 Map
000	No regional lymph node involvement	N0	N0	NONE	NONE

Code	Description	TNM 7 Map	TNM 6 Map	SS77 Map	SS2000 Map
100	Regional lymph node(s): Iliac, NOS: Common External Internal (hypogastric) Obturator Pelvic, NOS	N1	N1	RN	RN
120	OBSOLETE DATA RETAINED V0200 Regional lymph node(s): Lateral sacral (laterosacral) Presacral	ERROR	N1	D	RN
200	Regional lymph node(s): Aortic, NOS: Lateral (lumbar) Para-aortic Periaortic Retroperitoneal, NOS	N1	N1	RN	RN
220	OBSOLETE DATA RETAINED V0200 20 + 12	ERROR	N1	D	RN
300	Regional lymph node(s): Inguinal	N1	N1	D	RN
500	Regional lymph node(s), NOS	N1	N1	RN	RN
800	Lymph nodes, NOS	N1	N1	RN	RN
999	Unknown; not stated Regional lymph node(s) cannot be assessed Not documented in patient record	NX	NX	U	U

FallopianTube**Reg LN Pos****Note:** Record this field even if there has been preoperative treatment.**See Standard Table****FallopianTube****Reg LN Exam****See Standard Table****FallopianTube****CS Mets at DX**

Code	Description	TNM 7 Map	TNM 6 Map	SS77 Map	SS2000 Map
00	No; none	M0	M0	NONE	NONE
05	Presacral lymph nodes Lateral sacral (laterosacral) lymph nodes Sacral NOS	M1	*	D	RN
10	Distant lymph node(s), NOS other than presacral	M1	M1	D	D
20	10 + 05	M1	*	D	D
40	Distant metastases, except distant lymph nodes (code 10), including: Liver parenchymal metastasis Pleural effusion WITH positive cytology Carcinomatosis	M1	M1	D	D
45	40 + 05	M1	*	D	D
50	40 + 10 Distant lymph node(s) other than presacral plus other distant metastases	M1	M1	D	D
55	50 + 05	M1	*	D	D

Code	Description	TNM 7 Map	TNM 6 Map	SS77 Map	SS2000 Map
60	Distant metastasis, NOS Stated as M1 [NOS]	M1	M1	D	D
99	Unknown if distant metastasis Distant metastasis cannot be assessed Not documented in patient record	M0	MX	U	U

* For codes 05, 20, 45, and 55, the N and M categories are assigned based on the coding of this CS Mets at DX and CS Lymph Nodes - see table AJCC6th N and M from CS Lymph nodes and CS Mets at DX.