

**Scrotum [Merkel Cell Carcinoma only]****C63.2**

(M-8247)

C63.2 Scrotum, NOS

**MerkelCellScrotum****CS Tumor Size**

**Note:** Code the specific tumor size as documented in the medical record. If the ONLY information regarding tumor size is the physician's statement of the T category, assign code 992, 995 or 996.

<b>Code</b>	<b>Description</b>
000	No mass/tumor found
001-988	001 - 988 millimeters (code exact size in millimeters)
989	989 millimeters or larger
990	Microscopic focus or foci only and no size of focus given
991	Described as "less than 1 cm"
992	Described as "less than 2 cm," or "greater than 1 cm," or "between 1 cm and 2 cm" Stated as T1 with no other information on size
993	Described as "less than 3 cm," or "greater than 2 cm," or "between 2 cm and 3 cm"
994	Described as "less than 4 cm," or "greater than 3 cm," or "between 3 cm and 4 cm"
995	Described as "less than 5 cm," or "greater than 4 cm," or "between 4 cm and 5 cm" Stated as T2 with no other information on size
996	Described as "greater than 5 cm" Stated as T3 with no other information on size
999	Unknown; size not stated Not documented in patient record

**MerkelCellScrotum****CS Extension**

**Note 1:** In the case of multiple simultaneous tumors, code the tumor with greatest extension.

**Note 2:** Skin ulceration does not alter the Collaborative Stage classification.

**Note 3:** In-transit metastasis is coded in the CS Lymph Nodes field.

**Note 4:** If the ONLY information regarding the tumor extension is the physician's statement of the T category, assign code 310, 320, 330, or 810. (Refer to CS Tumor Size table for instructions on coding tumor size).

Code	Description	TNM 7 Map	TNM 6 Map	SS77 Map	SS2000 Map
000	In situ; noninvasive; intraepidermal	Tis	Tis	IS	IS
100	Lesion(s) confined to dermis Confined to scrotum	^	*	L	L
300	Localized, NOS	^	*	L	L
305	Subcutaneous tissue (through entire dermis)	^	T1	L	L
310	Stated as T1 with no other information on extension	^	*	L	L
320	Stated as T2 with no other information on extension	^	T2	L	L
330	Stated as T3 with no other information on extension	^	T3	L	L
400	Adjacent connective tissue (See definition of connective tissue in general instructions)	^	*	RE	RE
600	Adjacent organs/structures Male genital organs: Epididymis Penis Prostate Spermatic cord Testis	^	T4	RE	RE

Code	Description	TNM 7 Map	TNM 6 Map	SS77 Map	SS2000 Map
800	Further contiguous extension Other organs and structures in male pelvis: Bladder Rectum Urethra	^	T4	D	D
805	Underlying cartilage, bone, skeletal muscle	T4	T4	D	D
810	Stated as T4 with no other information on extension	T4	T4	RE	RE
950	No evidence of primary tumor	T0	T0	U	U
999	Unknown extension Primary tumor cannot be assessed Not documented in patient record	TX	TX	U	U

^ For CS Extension codes 100, 300, 305, 310, 320, 330, 400, 600 and 800 ONLY, the T category for AJCC 7th Edition is assigned based on the value of CS Tumor Size and CS Extension as shown in the Extension Size Table AJCC 7 for this site.

\* For CS Extension codes 100, 300, 310 and 400 ONLY, the T category for AJCC 6th Edition is assigned based on the value of CS Tumor Size, as shown in the Extension Size Table AJCC 6 for this site.

### **MerkelCellScrotum**

#### **CS Tumor Size/Ext Eval**

See Standard Table

### **MerkelCellScrotum**

#### **CS Lymph Nodes**

**Note 1:** Code only regional nodes and nodes, NOS, in this field. Distant nodes are coded in the field Mets at DX.

**Note 2:** In-transit metastasis is defined as a tumor distinct from the primary lesion and located either between the primary lesion and the draining regional lymph nodes or distal to the primary lesion. In-transit metastasis is coded in the CS Lymph Nodes field.

**Note 3:** According to AJCC, "there is no lower threshold of tumor burden defining the presence of regional nodal metastasis. Specifically, nodal tumor deposits less than 0.2 mm in diameter (previously used as the threshold for defining nodal metastasis), are included in the staging of nodal disease as a result of the consensus that smaller volumes of metastatic tumor are still clinically significant". This

means that any finding of Merkel cell carcinoma in lymph nodes, regardless of size and including isolated tumor cells (ITCs), should be coded positive lymph nodes in this field.

Code	Description	TNM 7 Map	TNM 6 Map	SS77 Map	SS2000 Map
000	None; no regional lymph node involvement	N0	N0	NONE	NONE
100	Regional lymph nodes Iliac, NOS: External Internal (hypogastric), NOS: Obturator Inguinal, NOS: Deep inguinal, NOS Node of Cloquet or Rosenmuller (highest deep inguinal) Superficial inguinal (femoral) Regional lymph node(s), NOS	^	N1	RN	RN
320	Stated as N1a with no other information on regional lymph nodes	N1a	N1	RN	RN
340	Stated as N1b with no other information on regional lymph nodes	N1b	N1	RN	RN
360	Stated as N1 [NOS] with no other information on regional lymph nodes	N1NOS	N1	RN	RN
400	In-transit metastasis WITHOUT regional lymph node involvement or involvement of regional lymph nodes not stated (see Note 2)	N2	N1	RN	RN
420	400 + 100 In-transit metastasis WITH regional lymph nodes listed in code 100 (see Note 2)	N2	N1	RN	RN
800	Lymph nodes, NOS	^	N1	RN	RN
999	Unknown; not stated	NX	NX	U	U

Code	Description	TNM 7 Map	TNM 6 Map	SS77 Map	SS2000 Map
999 cont'd	Regional lymph node(s) cannot be assessed Not documented in patient record	NX	NX	U	U

^ For codes 100, 200, 300, 400, 500 and 800 ONLY, the N category for AJCC 7th Edition is assigned based on the value of Site-Specific Factor 3, Clinical Status of Lymph Node Mets, as shown in the CS Lymph Nodes and Clinical Status table.

### MerkelCellScrotum

#### Reg LN Pos

**Note:** Record this field even if there has been preoperative treatment.

**See Standard Table**

### MerkelCellScrotum

#### Reg LN Exam

**See Standard Table**

### MerkelCellScrotum

#### CS Mets at DX

**Note:** In-transit metastasis is coded in the CS Lymph Nodes field.

Code	Description	TNM 7 Map	TNM 6 Map	SS77 Map	SS2000 Map
00	No; none	M0	M0	NONE	NONE
10	Distant lymph node(s)	M1a	M1	D	D
15	Metastasis to skin or subcutaneous tissue	M1a	M1	D	D
20	Stated as M1a with no other information on distant metastases	M1a	M1	D	D
30	Lung	M1b	M1	D	D
35	30 + (10 or 15) Metastasis to lung plus distant lymph node(s) or skin or subcutaneous tissue	M1b	M1	D	D
37	Stated as M1b with no other	M1b	M1	D	D

Code	Description	TNM 7 Map	TNM 6 Map	SS77 Map	SS2000 Map
37 cont'd	information on distant metastases	M1b	M1	D	D
40	OBSOLETE DATA RETAINED V0200 Distant metastasis, NOS Distant metastases except distant lymph node(s) (code 10) Carcinomatosis	ERROR	M1	D	D
45	All other visceral sites Distant metastases except distant lymph node(s), skin or subcutaneous tissue, or lung (codes 10, 15, 30 or 35) Carcinomatosis	M1c	M1	D	D
50	OBSOLETE DATA RETAINED V0200 (10) + (40) Distant lymph node(s) plus other distant metastases	ERROR	M1	D	D
55	45 + (any of 10, 15 or 30)Metastases to other visceral sites plus any of distant lymph node(s), skin, subcutaneous tissue, or lung	M1c	M1	D	D
57	Stated as M1c with no other information on distant metastases	M1c	M1	D	D
60	Distant metastasis, NOS Stated as M1 [NOS] with no other information on distant metastases	M1NOS	M1	D	D
99	Unknown if distant metastasis Distant metastasis cannot be assessed Not documented in patient record	M0	MX	U	U