

Larynx Supraglottic
Supraglottic Larynx (excluding Malignant Melanoma)

C32.1

C32.1 Supraglottis

Note: Excludes Anterior Surface of Epiglottis - see separate schema (C10.1).

Larynx Supraglottic

CS Tumor Size

See Standard Table

Larynx Supraglottic

CS Extension

Note 1: Impaired vocal cord mobility may be described as vocal cord paresis, and may suggest invasion of intrinsic laryngeal muscle. Fixation of the vocal cord may be described as immobility of the arytenoids noted on endoscopy, paralysis of the vocal cords, or deviation of larynx to the fixed side.

Note 2: Use code 450 for localized tumor only if no information is available to assign codes 100 through 400.

Note 3: Use code 685, 735, 810, or 815 if the physician's assignment of T category is the only information available about the extent of the tumor.

Code	Description	TNM 7 Map	TNM 6 Map	SS77 Map	SS2000 Map
000	In situ, intraepithelial, noninvasive	Tis	Tis	IS	IS
100	Invasive tumor confined to: Supraglottis (one subsite): Aryepiglottic fold Arytenoid cartilage Corniculate cartilage Cuneiform cartilage Epilarynx, NOS False cords: Ventricular bands Ventricular cavity Ventricular fold Infrahyoid epiglottis Laryngeal cartilage, NOS Laryngeal (posterior) surface of epiglottis Suprahyoid epiglottis (including tip, lingual (anterior) and laryngeal surfaces) WITH normal vocal cord	T1	T1	L	L

Code	Description	TNM 7 Map	TNM 6 Map	SS77 Map	SS2000 Map
100 cont'd	mobility Stated as T1 with no further information on extension	T1	T1	L	L
200	Tumor involves mucosa of more than one subsite of supraglottis listed in code 100 WITHOUT impaired vocal cord mobility or fixation of larynx or NOS	T2	T2	L	L
230	Tumor involves mucosa of glottis WITHOUT impaired vocal cord mobility or fixation of larynx or NOS	T2	T2	L	L
240	Tumor involved adjacent subsites in supraglottis or glottis, NOS, mucosal involvement not specified, WITHOUT impaired vocal cord mobility or fixation of larynx or NOS	T2	T2	L	L
250	Involvement of any structures in supraglottic larynx listed in code 100 and/or glottis WITH impaired vocal cord mobility	T2	T2	L	L
300	OBSOLETE DATA RETAINED V0200 See code 240 Tumor involves adjacent region(s) of larynx	ERROR	T2	L	L
320	Tumor involves mucosa of regions outside supraglottis, including: Base of tongue Vallecula Medial wall of pyriform sinus WITHOUT impaired vocal cord mobility or fixation of larynx or NOS	T2	T2	RE	RE

Code	Description	TNM 7 Map	TNM 6 Map	SS77 Map	SS2000 Map
350	OBSOLETE DATA RETAINED V0200 See codes 250 and 360 Impaired vocal cord mobility	T2	T2	L	L
360	Involvement of any structures in code 320 WITH impaired vocal cord mobility	T2	T2	RE	RE
365	Tumor involves adjacent regions of supraglottis or glottis or region outside supraglottis, NOS, mucosal involvement not specified, WITHOUT impaired vocal cord mobility or fixation of larynx or NOS	T2	T2	RE	RE
375	Stated as T2 with no other information on extension	T2	T2	L	L
390	Involvement of subglottis WITHOUT vocal cord fixation	T3	T3	L	L
395	390 + 320 Subglottis plus any structure in code 320 WITHOUT vocal cord fixation	T3	T3	RE	RE
400	Involvement of any structures in supraglottis (code 100), glottis, and subglottis WITH vocal cord fixation	T3	T3	L	L
450	Localized, NOS	T1	T1	L	L
520	Paraglottic space WITHOUT vocal cord fixation	T3	T3	RE	RE
530	520 + 320	T3	T3	RE	RE
600	OBSOLETE DATA RETAINED V0200 See code 320	ERROR	T2	RE	RE

Code	Description	TNM 7 Map	TNM 6 Map	SS77 Map	SS2000 Map
600 cont'd	Tumor involves region outside the supraglottis WITHOUT fixation, including: Medial wall pyriform sinus Mucosa of base of tongue Vallecula	ERROR	T2	RE	RE
620	OBSOLETE DATA RETAINED V0200 Code 600 WITH fixation	ERROR	T3	RE	RE
625	Involvement of any structures in code 320 and/or 520 WITH vocal cord fixation	T3	T3	RE	RE
650	Hypopharynx, NOS Postcricoid area Pre-epiglottic tissues	T3	T3	RE	RE
660	Deep base of tongue	T3	T3	RE	RE
670	OBSOLETE DATA CONVERTED V0200; See code 690 Cricoid cartilage	ERROR	ERROR	ERROR	ERROR
680	Inner cortex of thyroid cartilage (minor erosion)	T3	T3	RE	D
685	Stated as T3 with no other information on extension	T3	T3	RE	RE
690	Cricoid cartilage	T4a	T4a	RE	RE
695	690 + 680 Cricoid cartilage plus inner cortex of thyroid cartilage	T4a	T4a	RE	D
700	Extension to/through: Esophagus Oropharynx	T4a	T4a	D	D

Code	Description	TNM 7 Map	TNM 6 Map	SS77 Map	SS2000 Map
700 cont'd	Soft tissues of neck Outer cortex of thyroid cartilage Thyroid gland	T4a	T4a	D	D
720	Extension to/through: Strap muscle(s) Omohyoid Sternohyoid Sternothyroid Thyrohyoid Skin	T4a	T4a	D	D
730	Extension to/through: Deep extrinsic muscle(s) of tongue Trachea	T4a	T4a	D	D
735	Stated as T4a with no other information on extension	T4a	T4a	RE	RE
800	Further contiguous extension, including: Carotid artery (encased) Mediastinal structures Prevertebral space	T4b	T4b	D	D
810	Stated as T4b with no other information on extension	T4b	T4b	D	D
815	Stated as T4 [NOS] with no other information on extension	T4NOS	T4NOS	RE	RE
950	No evidence of primary tumor	T0	T0	U	U
999	Unknown; extension not stated Primary tumor cannot be assessed Not documented in patient record	TX	TX	U	U

**Larynx Supraglottic
CS Tumor Size/Ext Eval
See Standard Table**

**Larynx Supraglottic
CS Lymph Nodes**

Note 1: For head and neck schemas, this field includes all lymph nodes defined as Levels I-VII and Other by AJCC. The complete definitions are provided in the General Instructions.

Note 2: For head and neck schemas, additional information about lymph nodes (size of involved nodes, extracapsular extension, levels involved, and location of involved nodes above or below the lower border of the cricoid cartilage) is coded in Site-Specific Factors 1, 3-9.

Note 3: If laterality of lymph nodes is not specified, assume nodes are ipsilateral. Midline nodes are considered ipsilateral.

Note 4: For head and neck cancers, if lymph nodes are described only as "supraclavicular", try to determine if they are in Level IV (deep to the sternocleidomastoid muscle, in the lower jugular chain) or Level V (in the posterior triangle, inferior to the transverse cervical artery) and code appropriately. If the specific level cannot be determined, consider them as Level V nodes.

Note 5: The description of lymph nodes has been standardized across the head and neck schemas. All lymph node levels and groups listed here are considered regional nodes for AJCC staging.

Summary Stage 1977 and Summary Stage 2000 divide these nodes into regional and distant groups.

Note 6: Level III nodes have been moved from code 100 in CSV1 to code 110. Level IV nodes have been added to code 120.

^ For codes 100-120, 200-220, 300-320, 400-420, 500-520, and 800 ONLY, the N category for AJCC 7th Edition staging is assigned based on the value of Site-Specific Factor 1, Size of Lymph Nodes, using the extra table Lymph Nodes Size Table, for this site.

* For codes 100-120, 200-220, 300-320, 400-420, 500-520, and 800 ONLY, the N category for AJCC 6th Edition staging is assigned based on the value of Site-Specific Factor 1, Size of Lymph Nodes, using the extra table Lymph Nodes Size Table, for this site.

Code	Description	TNM 7 Map	TNM 6 Map	SS77 Map	SS2000 Map
000	No regional lymph node involvement	N0	N0	NONE	NONE
100	Single positive ipsilateral regional node: Level II - Upper jugular: Jugulodigastric (subdigastric) Upper deep cervical Level IIA - Anterior Level IIB - Posterior Level VI node - Anterior compartment group: Laterotracheal Paralaryngeal	^	*	RN	RN

Code	Description	TNM 7 Map	TNM 6 Map	SS77 Map	SS2000 Map
100 cont'd	Paratracheal - above suprasternal notch Perithyroidal Precricoid (Delphian) Pretracheal - above suprasternal notch Recurrent laryngeal Cervical, NOS Deep cervical, NOS Internal jugular NOS: Regional lymph node, NOS	^	*	RN	RN
110	Single positive ipsilateral regional node: Level I: Level IA - Submental Level IB - Submandibular (submaxillary), sublingual Level III - Middle jugular: Middle deep cervical Retropharyngeal	^	*	D	RN
120	Single positive ipsilateral regional node: Level IV - Lower jugular Jugulo-omohyoid (supraomohyoid) Lower deep cervical Virchow node Level V - Posterior triangle group: Level VA - Spinal accessory Level VB - Transverse cervical, supraclavicular (see Note 4) Level VII node - Superior mediastinal group (for other mediastinal nodes see CS Mets at DX): Esophageal groove Paratracheal - below suprasternal notch Pretracheal - below suprasternal notch Other groups: Facial: Buccinator (buccal) Mandibular Nasolabial Parotid:	^	*	D	D

Code	Description	TNM 7 Map	TNM 6 Map	SS77 Map	SS2000 Map
120 cont'd	Infraauricular Intraparotid Periparotid Preparotid Parapharyngeal Retroauricular (mastoid) Suboccipital	^	*	D	D
180	Stated as N1 with no other information on regional lymph nodes	^	*	RN	RN
190	Stated as N2a with no other information on regional lymph nodes	^	*	RN	RN
200	Multiple positive ipsilateral nodes listed in code 100	^	*	RN	RN
210	Multiple positive ipsilateral nodes, any listed in code 110 WITH or WITHOUT nodes listed in code 100	^	*	D	RN
220	Multiple positive ipsilateral nodes, any listed in code 120 WITH or WITHOUT nodes listed in code 100 or 110	^	*	D	D
290	Stated as N2b with no other information on regional lymph nodes	^	*	RN	RN
300	Positive ipsilateral node(s) listed in code 100, not stated if single or multiple	^	*	RN	RN
310	Positive ipsilateral node(s) listed in code 110, not stated if single or multiple	^	*	D	RN
320	Positive ipsilateral node(s) listed in code 120, not stated if single or multiple	^	*	D	D
400	Positive bilateral or contralateral nodes listed in code 100	^	*	RN	RN

Code	Description	TNM 7 Map	TNM 6 Map	SS77 Map	SS2000 Map
410	Positive bilateral or contralateral nodes, any listed in code 110 WITH or WITHOUT nodes listed in code 100	^	*	D	RN
420	Positive bilateral or contralateral nodes, any listed in code 120 WITH or WITHOUT nodes listed in code 100 or 110	^	*	D	D
490	Stated as N2c with no other information on regional lymph nodes	^	*	RN	RN
500	Positive node(s) listed in code 100, not stated if ipsilateral, or bilateral, or contralateral, AND not stated if single or multiple	^	*	RN	RN
510	Positive node(s) listed in code 110, not stated if ipsilateral, or bilateral, or contralateral, AND not stated if single or multiple	^	*	D	RN
520	Positive node(s) listed in code 120, not stated if ipsilateral, or bilateral, or contralateral, AND not stated if single or multiple	^	*	D	D
600	Stated as N2 [NOS] with no other information on regional lymph nodes	^	*	RN	RN
700	Stated as N3 with no other information on regional lymph nodes	^	*	RN	RN
800	Lymph nodes, NOS	N1	N1	RN	RN
999	Unknown; regional lymph nodes not stated Regional lymph nodes cannot be assessed	NX	NX	U	U

Code	Description	TNM 7 Map	TNM 6 Map	SS77 Map	SS2000 Map
999 cont'd	Not documented in patient record	NX	NX	U	U

^ For codes 100-120, 200-220, 300-320, 400-420, 500-520, and 800 ONLY, the N category for AJCC 7th Edition staging is assigned based on the value of Site-Specific Factor 1, Size of Lymph Nodes, using the extra table Lymph Nodes Size Table, for this site.

* For codes 100-120, 200-220, 300-320, 400-420, 500-520, and 800 ONLY, the N category for AJCC 6th Edition staging is assigned based on the value of Site-Specific Factor 1, Size of Lymph Nodes, using the extra table Lymph Nodes Size Table, for this site.

**Larynx Supraglottic
CS Lymph Nodes Eval
See Standard Table**

**Larynx Supraglottic
Regional Nodes Positive
See Standard Table**

Note: Record this field even if there has been preoperative treatment.

**Larynx Supraglottic
Regional Nodes Examined
See Standard Table**

**Larynx Supraglottic
CS Mets at DX**

Note: Supraclavicular and transverse cervical lymph nodes are coded in CS Lymph Nodes because they are categorized as N rather than M in AJCC TNM.

Code	Description	TNM 7 Map	TNM 6 Map	SS77 Map	SS2000 Map
00	No distant metastasis	M0	M0	NONE	NONE
10	Distant lymph node(s): Mediastinal (excluding superior mediastinal nodes) Distant lymph node(s), NOS	M1	M1	D	D
40	Distant metastases except distant lymph node(s) Carcinomatosis	M1	M1	D	D

Code	Description	TNM 7 Map	TNM 6 Map	SS77 Map	SS2000 Map
50	40+10 Distant metastasis plus distant lymph node(s)	M1	M1	D	D
60	Distant metastasis, NOS Stated as M1 with no other information on distant metastasis	M1	M1	D	D
99	Unknown; distant metastasis not stated Distant metastasis cannot be assessed Not documented in patient record	M0	MX	U	U

Larynx Supraglottic**CS Mets Eval****See Standard Table**

Note: This item reflects the validity of the classification of the item CS Mets at DX only according to the diagnostic methods employed.

Larynx Supraglottic**CS Site-Specific Factor 1****Size of Lymph Nodes****Note: See page A-96**

Note: Code the largest diameter, whether measured clinically or pathologically, of any involved regional lymph node(s). Do not code the size of any nodes coded in CS Mets at DX.

Code	Description
000	No involved regional nodes
001-979	001 - 979 millimeters (mm); (Exact size of lymph node to nearest mm)
980	980 mm or larger (Includes cases converted from codes 981-989 during conversion to V0200)
981-987	OBSOLETE DATA CONVERTED V0200 See code 980 981 - 987 mms
988	Not applicable: Information not collected for this case (If this item is required by your standard setter, use of code 988 will result in an

Code	Description
988 cont'd	edit error)
989	OBSOLETE DATA CONVERTED V0200 See code 980 989 mm or larger
990	Microscopic focus or foci only and no size of focus given
991	Described as "less than 1centimeter (cm)"
992	Described as "less than 2cm" or "greater than 1cm" or "between 1cm and 2cm"
993	Described as "less than 3cm" or "greater than 2cm" or "between 2cm and 3cm"
994	Described as "less than 4cm" or "greater than 3cm" or "between 3cm and 4cm"
995	Described as "less than 5cm" or "greater than 4cm" or "between 4cm and 5cm"
996	Described as "less than 6cm" or "greater than 5cm" or "between 5cm and 6cm"
997	Described as "more than 6cm"
999	Regional lymph node(s) involved, size not stated Unknown if regional lymph nodes involved Not documented in patient record