

Fallopian Tube**C57.0****C57.0 Fallopian tube**

Note 1: Laterality must be coded for this site.

Note 2: AJCC TNM values correspond to the stages accepted by the Federation Internationale de Gynecologic et d'Obstetrique (FIGO). Note that FIGO uses a single stage value which corresponds to different groupings of T, N, and M values.

FallopianTube**CS Tumor Size**

See Standard Table

FallopianTube**CS Extension**

Note 1: When both Federation Internationale de Gynecologie et d'Obstetrique (FIGO) stage and extension detail are available, record the code with extension detail in preference to a statement of FIGO stage. FIGO Stage IIIC is assigned based on tumor extension and/or regional lymph node involvement. Code CS Extension based on a physician's statement of FIGO Stage IIIC only if it is known that the stage value is assigned because of tumor extension.

Note 2: FIGO no longer includes a Stage 0 (Tis) for this schema.

Note 3: Liver capsule metastases are coded in CS Extension; liver parenchymal metastases are coded in CS Mets at DX.

Note 4: Both extension to and discontinuous metastasis to any of the following pelvic organs are considered FIGO Stage II and coded in the range 400-660: adnexa, NOS; bladder, bladder serosa; broad ligament (mesovarium); cul de sac; contralateral fallopian tube; parametrium; pelvic peritoneum; pelvic wall; rectum; sigmoid colon; sigmoid mesentery; ureter; uterus; uterine serosa.

Note 5: Both extension to and discontinuous metastasis to any of the following abdominal organs are considered FIGO Stage III and coded in the range 685-795: abdominal mesentery; diaphragm; gallbladder; infracolic omentum; kidneys; large intestine except rectum and sigmoid colon; liver (peritoneal surface); omentum; pancreas; pericolic gutter; peritoneum, NOS; small intestine; spleen; stomach; ureters.

Note 6: From the AJCC Manual 7th Edition (page 430): "It may be preferable to classify a patient as TX (primary tumor cannot be assessed) if inadequate staging biopsies and/or a lack of peritoneal cytology make it inaccurate to classify the patient with confidence as early stage (Stage T3a/IIIA has not been excluded by adequate staging biopsies)."

Note 7: Since "malignant ascites or malignant peritoneal washings" was not specifically categorized in the 1977 Summary Staging Guide, it is unclear to which stage pre-2001 cases may have been coded.

Code	Description	TNM 7 Map	TNM 6 Map	SS77 Map	SS2000 Map
000	In situ, intraepithelial, noninvasive Limited to tubal mucosa	Tis	Tis	IS	IS

Code	Description	TNM 7 Map	TNM 6 Map	SS77 Map	SS2000 Map
100	OBSOLETE DATA RETAINED AND REVIEWED V0203 See codes 200, 320 Confined to fallopian tube, NOS FIGO Stage I	T1NOS	T1NOS	L	L
110	Confined to one fallopian tube WITHOUT penetrating serosal surface WITHOUT ascites FIGO Stage IA	T1a	T1a	L	L
120	Confined to both fallopian tubes WITHOUT penetrating serosal surface WITHOUT ascites FIGO Stage IB	T1b	T1b	L	L
130	Extension onto or through tubal serosa Malignant ascites Malignant peritoneal washings FIGO Stage IC	T1c	T1c	L	L
200	Confined to fallopian tube, NOS	T1NOS	T1NOS	L	L
300	OBSOLETE DATA RETAINED AND REVIEWED V0203 See codes 310, 320 Localized, NOS FIGO Stage I	T1NOS	T1NOS	L	L
310	Localized, NOS	T1NOS	T1NOS	L	L
320	FIGO Stage I [NOS]	T1NOS	T1NOS	L	L
350	OBSOLETE DATA RETAINED AND REVIEWED V0203 See code 655, 670 Pelvic extension, NOS with no malignant cells in ascites or peritoneal washings	T2NOS	T2NOS	RE	RE

Code	Description	TNM 7 Map	TNM 6 Map	SS77 Map	SS2000 Map
350 cont'd	FIGO Stage II	T2NOS	T2NOS	RE	RE
400	Extension or metastasis to: Corpus uteri Ovary, ipsilateral Uterus, NOS WITHOUT malignant cells in ascites or peritoneal washings	T2a	T2a	RE	RE
450	Extension or metastasis to: Ovary, contralateral WITHOUT malignant cells in ascites or peritoneal washings	T2a	T2a	D	RE
460	FIGO Stage IIA	T2a	T2a	RE	RE
500	OBSOLETE DATA RETAINED AND REVIEWED V0203 See codes 510, 620, and 658 Extension or metastasis to (but no malignant cells in ascites or peritoneal washings): Broad ligament, ipsilateral Mesosalpinx, ipsilateral Adjacent peritoneum FIGO Stage IIB	T2b	T2b	RE	RE
510	Extension or metastasis to: Broad ligament, ipsilateral Mesosalpinx, ipsilateral Adjacent peritoneum WITHOUT malignant cells in ascites or peritoneal washings	T2b	T2b	RE	RE
600	OBSOLETE DATA RETAINED V0200; See codes 450, 460 Ovary, contralateral (but no malignant cells in ascites or peritoneal washings) FIGO Stage IIA	ERROR	T2a	D	RE

Code	Description	TNM 7 Map	TNM 6 Map	SS77 Map	SS2000 Map
610	OBSOLETE DATA RETAINED V0200 (600) + (500)	ERROR	T2b	D	RE
620	51 0 + 450 Extension to any site in 500 plus extension to contralateral ovary WITHOUT malignant cells in ascites or peritoneal washings	T2b	T2b	D	RE
650	OBSOLETE DATA RETAINED AND REVIEWED V0203 See codes 652, 658 Extension or metastasis to (but no malignant cells in ascites or peritoneal washings): Broad ligament, contralateral Cul de sac (rectouterine pouch) Mesosalpinx, contralateral Rectosigmoid Sigmoid FIGO IIB	T2b	T2b	D	RE
652	Extension or metastasis to (but no malignant cells in ascites or peritoneal washings): Broad ligament, contralateral Cul de sac (rectouterine pouch) Mesosalpinx, contralateral Rectosigmoid Sigmoid	T2b	T2b	D	RE
655	Pelvic extension, NOS with no malignant cells in ascites or peritoneal washings	T2NOS	T2NOS	RE	RE
658	FIGO Stage IIB	T2b	T2b	RE	RE
660	Pelvic extension (codes 400-655)	T2c	T2c	D	D

Code	Description	TNM 7 Map	TNM 6 Map	SS77 Map	SS2000 Map
660 cont'd	WITH malignant cells in ascites or peritoneal washings FIGO Stage IIC	T2c	T2c	D	D
670	FIGO Stage II [NOS]	T2NOS	T2NOS	RE	RE
680	OBSOLETE DATA RETAINED AND REVIEWED V0203 See code 790, 820 Peritoneal implants or metastasis (size of metastases not stated; unknown if microscopic or macroscopic) Omentum Small intestine FIGO Stage III	T3NOS	T3NOS	D	RE
685	Microscopic peritoneal implants or metastasis: Omentum Small intestine	T3a	T3a	D	RE
688	685 + 660	T3a	T3a	D	D
690	OBSOLETE DATA RETAINED AND REVIEWED V0203 See codes 685, 698 Microscopic peritoneal implants or metastasis: Omentum Small intestine FIGO Stage IIIA	T3a	T3a	D	RE
695	Microscopic peritoneal metastasis outside the pelvis, excluding organs in code 685 (See Note 5)	T3a	T3a	D	D
698	FIGO Stage IIIA	T3a	T3a	D	RE

Code	Description	TNM 7 Map	TNM 6 Map	SS77 Map	SS2000 Map
700	OBSOLETE DATA CONVERTED V0102 ; See code 680 Extension or metastasis to: Omentum FIGO Stage IIB	ERROR	ERROR	ERROR	ERROR
710	OBSOLETE DATA CONVERTED V0102 ; See code 660 Pelvic extension (codes 350-700) with malignant cells in ascites or peritoneal washings FIGO Stage IIC	ERROR	ERROR	ERROR	ERROR
720	OBSOLETE DATA RETAINED AND REVIEWED V0203 See codes 722, 723, and 728 Macroscopic peritoneal implants or metastasis less than or equal to 2cm: Omentum Small intestine FIGO Stage IIIB	T3b	T3b	D	RE
722	Macroscopic peritoneal implants or metastasis less than or equal to 2cm: Omentum Small intestine	T3b	T3b	D	RE
723	722 + 695	T3b	T3b	D	D
725	Macroscopic peritoneal metastasis less than or equal to 2 cm outside the pelvis, excluding organs in code 722 (See Note 5)	T3b	T3b	D	D
728	FIGO Stage IIIB based on extension	T3b	T3b	D	RE
730	OBSOLETE DATA RETAINED AND REVIEWED V0203 See codes 732, 733, and 740 Macroscopic peritoneal implants or	T3c	T3c	D	RE

Code	Description	TNM 7 Map	TNM 6 Map	SS77 Map	SS2000 Map
730 cont'd	metastasis greater than 2cm: Omentum Small intestine FIGO Stage IIIC	T3c	T3c	D	RE
732	Macroscopic peritoneal implants or metastasis greater than 2cm: Omentum Small intestine	T3c	T3c	D	RE
733	732 + 725	T3c	T3c	D	D
735	Macroscopic peritoneal implants or metastases greater than 2 cm outside the pelvis, excluding organs in code 732 (See Note 5)	T3c	T3c	D	D
740	FIGO Stage IIIC based on extension	T3c	T3c	D	RE
750	OBSOLETE DATA CONVERTED V0203; See code 795 Peritoneal implants outside the pelvis, NOS (size of metastases not stated; unknown if microscopic or macroscopic), except code 680 (See Note 5)	ERROR	ERROR	ERROR	ERROR
760	OBSOLETE DATA RETAINED AND REVIEWED V0203 See codes 695 and 698 Microscopic peritoneal metastasis outside the pelvis, except code 690 (See Note 5) FIGO Stage IIIA	T3a	T3a	D	D
770	OBSOLETE DATA RETAINED AND REVIEWED V0203 See codes 725 and 728 Macroscopic peritoneal metastasis less	T3b	T3b	D	D

Code	Description	TNM 7 Map	TNM 6 Map	SS77 Map	SS2000 Map
770 cont'd	than or equal to 2 cm outside the pelvis, except code 720 (See Note 5) FIGO Stage IIIB	T3b	T3b	D	D
780	OBSOLETE DATA RETAINED AND REVIEWED V0200 See codes 735 and 740 Peritoneal metastases greater than 2 cm, except code 730 (See Note 5) FIGO Stage IIIC	T3c	T3c	D	D
790	Peritoneal implants or metastasis outside the pelvis, size of metastasis not stated, unknown if microscopic or macroscopic: Omentum Small intestine	T3NOS	T3NOS	D	RE
795	Peritoneal implants or metastases outside the pelvis, size of metastasis not stated, unknown if microscopic or macroscopic, excluding organs in code 790 (See Note 5)	T3NOS	T3NOS	D	D
800	OBSOLETE DATA RETAINED AND REVIEWED V0203 See codes 810, 820 Further contiguous extension FIGO Stage III [NOS]	T3NOS	T3NOS	D	D
810	Further contiguous extension	T3NOS	T3NOS	D	D
820	FIGO Stage III [NOS] based on extension	T3NOS	T3NOS	D	D
950	No evidence of primary tumor	T0	T0	U	U
999	Unknown; extension not stated Primary tumor cannot be assessed	TX	TX	U	U

Code	Description	TNM 7 Map	TNM 6 Map	SS77 Map	SS2000 Map
999 cont'd	(See Note 6) Not documented in patient record	TX	TX	U	U

Fallopian Tube**CS Tumor Size/Ext Eval**

See Standard Table

Fallopian Tube**CS Lymph Nodes**

Note 1: Code only regional nodes and nodes, NOS, in this field. Regional nodes include bilateral and contralateral involvement of named nodes. Distant nodes, including presacral and lateral sacral (laterosacral) nodes, are coded in CS Mets at DX.

Note 2: The Federation Internationale de Gynecologie et d'Obstetrique (FIGO) Stage IIIC is assigned based on tumor extension and/or regional lymph node involvement. Code CS Lymph Nodes based on a physician's statement of FIGO Stage IIIC if the stage value is assigned because of regional lymph node involvement or there is no indication of the basis.

Note 3: If either exploratory or definitive surgery is done with no mention of lymph nodes, assume nodes are negative.

Note 4: If the clinician says "adnexa palpated" but does not mention lymph nodes, assume lymph nodes are not involved.

Note 5: Determination of the N category for AJCC 6 is based on this field and CS Mets at DX for specific codes in CS Mets at DX.

Code	Description	TNM 7 Map	TNM 6 Map	SS77 Map	SS2000 Map
000	No regional lymph node involvement	N0	*	NONE	NONE
100	Regional lymph node(s): Iliac, NOS: Common External Internal (hypogastric) Obturator Pelvic, NOS	N1	N1	RN	RN
120	OBsolete DATA RETAINED V0200 Regional lymph node(s): Lateral sacral (laterosacral)	ERROR	N1	D	RN

Code	Description	TNM 7 Map	TNM 6 Map	SS77 Map	SS2000 Map
120 cont'd	Presacral	ERROR	N1	D	RN
200	Regional lymph node(s): Aortic, NOS: Lateral (lumbar) Para-aortic Periaortic Retroperitoneal, NOS	N1	N1	RN	RN
220	OBsolete DATA RETAINED V0200; 200 + 120	ERROR	N1	D	RN
300	Regional lymph node(s): Inguinal	N1	N1	D	RN
500	Regional lymph node(s), NOS	N1	N1	RN	RN
550	FIGO Stage IIIC based on regional lymph node involvement	N1	N1	RN	RN
800	Lymph nodes, NOS	N1	N1	RN	RN
999	Unknown; regional lymph nodes not stated Regional lymph node(s) cannot be assessed Not documented in patient record	NX	*	U	U

* For CS Lymph Nodes codes 000 and 999, the N category for AJCC 6 staging is based on the value of CS Mets at DX as shown in the Lymph Nodes Mets at DX AJCC 6 Table.

Fallopian Tube

CS Lymph Nodes Eval

See Standard Table

Fallopian Tube

Regional Nodes Positive

See Standard Table

Note: Record this field even if there has been preoperative treatment.

Fallopian Tube
Regional Nodes Examined
See Standard Table

Fallopian Tube
CS Mets at DX

Note 1: Discontinuous metastasis to any of the following abdominal organs by way of peritoneal seeding or implants is included in the T3 category (FIGO Stage III) and coded in CS Extension: abdominal mesentery; diaphragm; gallbladder; infracolic omentum; kidneys; large intestine except rectum, rectosigmoid, and sigmoid colon; liver (peritoneal surface); omentum; pancreas; pericolic gutter; peritoneum, NOS; small intestine; spleen; stomach; and ureters outside pelvis. Hematogenous metastases are included in M1 (FIGO Stage IV) disease.

Note 2: The Federation Internationale de Gynecologie et d'Obstetrique (FIGO) stages are included for coding where no more specific information is available in the record.

Code	Description	TNM 7 Map	TNM 6 Map	SS77 Map	SS2000 Map
00	No distant metastasis	M0	M0	NONE	NONE
05	Sacral lymph nodes: Presacral lymph nodes Lateral sacral (laterosacral) Sacral NOS	M1	M0	D	RN
10	Distant lymph node(s) other than sacral	M1	M1	D	D
20	10 + 05	M1	M1	D	D
40	Distant metastasis (excluding involvement of other organs by peritoneal seeding or implants (See Note 1)), except distant lymph nodes including: Liver parenchymal metastasis Pleural effusion WITH positive cytology Carcinomatosis	M1	M1	D	D
45	40 + 05	M1	M1	D	D
50	40 + 10 Distant metastasis plus distant lymph	M1	M1	D	D

Code	Description	TNM 7 Map	TNM 6 Map	SS77 Map	SS2000 Map
50 cont'd	node(s) other than sacral	M1	M1	D	D
55	50 + 05 Distant metastasis plus distant lymph nodes including sacral nodes	M1	M1	D	D
60	Distant metastasis, NOS FIGO Stage IV	M1	M1	D	D
99	Unknown; distant metastasis not stated Distant metastasis cannot be assessed Not documented in patient record	M0	MX	U	U

Fallopian Tube**CS Mets Eval****See Standard Table**

Note: This item reflects the validity of the classification of the item CS Mets at DX only according to the diagnostic methods employed.