

Appendix A. Denominators Used in Calculating Prevalence: Texas Resident Live Births, 1999-2013

Year	Breakdown		Texas Resident Live Births	
			Total	Males *
1999 through 2013, combined	Overall		5,740,466	2,934,020
	By Mother's Age	Less than 20 years	762,708	390,840
		20 to 24 years	1,571,886	801,595
		25 to 29 years	1,555,901	796,421
		30 to 34 years	1,189,596	607,459
		35 to 39 years	542,559	277,605
		40 or more years	117,343	59,872
		Unknown	473	228
	By Mother's Race/Ethnicity	White non-Hispanic	2,055,153	1,053,872
		Black non-Hispanic	645,316	328,671
		Hispanic	2,784,784	1,419,884
		Other non-Hispanic	248,237	127,971
		Unknown	6,976	3,622
	By Sex of Infant	Male	2,934,020	2,934,020
		Female	2,806,446	(NA)
	By Region of Mother's Residence at Delivery	Region 01	189,736	96,995
		Region 02	109,992	56,365
Region 03		1,532,305	784,861	
Region 04		218,154	111,720	
Region 05		151,381	77,428	
Region 06		1,390,462	709,564	
Region 07		625,736	319,911	
Region 08		562,391	286,806	
Region 09		131,879	67,556	
Region 10		216,835	110,372	
Region 11		611,595	312,442	
1999	Overall		349,157	178,451
2000	Overall		363,325	185,591
2001	Overall		365,092	186,774
2002	Overall		372,369	190,162
2003	Overall		377,374	192,581
2004	Overall		381,441	195,024
2005	Overall		385,537	197,491
2006	Overall		399,309	204,037
2007	Overall		407,453	208,222
2008	Overall		405,242	207,508
2009	Overall		401,599	204,876
2010	Overall		385,746	196,903
2011	Overall		377,274	192,766
2012	Overall		382,438	195,525
2013	Overall		387,110	198,109

* Male live births were the denominators for calculating the prevalence of hypospadias.