

# COVID-19 Vaccination Recommendations

January 7, 2022

	Primary Series			Booster Dose
	Dose 1	Dose 2	Dose 3 (Additional Dose) <sup>a</sup> (For Certain Immunocompromised Persons)	Single Booster Dose <sup>c</sup>
<b>Pfizer (5-11 years)</b> (orange cap)	0.2 mL <i>After dilution</i>	0.2 mL <sup>b</sup> @ 21 days after dose 1	0.2 mL <sup>b</sup> at least 28 days after dose 2	<b>NO</b> booster dose
<b>Pfizer (≥12 years)</b> purple → gray cap)	0.3 mL <i>(Gray cap does not require diluent)</i>	0.3 mL @ 21 days after dose 1	0.3 mL at least 28 days after dose 2	0.3 mL ≥ <b>5 Months</b> after dose 2 (primary series) or after additional dose <sup>a</sup>
<b>Moderna (≥18 years)</b>	0.5 mL	0.5 mL @ 28 Days after dose 1	0.5 mL at least 28 days after dose 2	0.25 mL ≥ <b>5 Months</b> after dose 2 (primary series) or after additional dose <sup>a</sup>
<b>J&amp;J's Janssen(≥18 years)<sup>d</sup></b>	0.5 mL	NA	NA	0.5 mL ≥ <b>2 Months</b> after dose 1

a: Certain Immunocompromised individuals only

b: Dose based on age at the time of the vaccination

c: Booster doses may be mix and match for individuals ≥18 yrs: [Booster dose may be mix-and-match vaccine](#)

d: [CDC preferentially recommends mRNA COVID-19 vaccines over Janssen COVID-19 vaccine](#)