


MATERNAL MORTALITY RATE: TEXAS AND THE UNITED STATES






Prepared by: Office of Program Decision Support, Division for Family and Community Health, Texas Department of State Health Services. Data Sources: Centers for Disease Control and Prevention, National Center for Health Statistics. Underlying Cause of Death and Natality public use data 2006-2015 on CDC Wonder Online Database. MMR computed within 42 days following the end of pregnancy, using ICD-10 codes A34, O00-O96, O98-O99.


MATERNAL MORTALITY RATES BY RACIAL/ETHNIC GROUP, 2006-2015



Prepared by: Office of Program Decision Support, Division for Family and Community Health Services, Texas Department of State Health Services, 07/21/2017. Data Source: Death and Birth Files, Center for Health Statistics, Texas Department of State Health Services. MMR - computed within 42 days following the end of pregnancy, using ICD-10 codes A34, O00-O95, O98-O99.



HISPANIC MOTHERS



Prepared by: Office of Program Decision Support, Division for Family and Community Health Services, Texas Department of State Health Services, 07/21/2017.

Data Sources: Death and Birth Files, Center for Health Statistics, Texas Department of State Health Services.

MMR—computed within 42 days following the end of pregnancy, using ICD-10 codes A34, O00-095, O98-O99.

Pre-pregnancy obesity; diabetes before and/or during pregnancy (including diabetes mellitus and gestational diabetes); and hypertension before and/or during pregnancy (including chronic hypertension, pre-eclampsia, and eclampsia)—Birth file. Note: r = correlation coefficient between MMR and each risk factor.