

Benefits and Risks of Vaccination

Vaccine-preventable disease	Effectiveness of vaccine	Possible vaccine side effects
Diphtheria - Bacterial toxins cause a thick covering in mucous membranes, most commonly in the back of the throat. It can lead to obstructive breathing problems, paralysis, heart failure, coma, and death within six-ten days.	Diphtheria toxoid is estimated to have a clinical efficacy of 97% against the disease.	DTaP/Tdap vaccine – Most children don't have any side effects from DTaP or Tdap. The side effects that do occur from DTaP or Tdap are usually mild, and may include, soreness or swelling at the injection site, fever, fussiness, feeling tired, loss of appetite or vomiting. Td should be used for those allergic to the pertussis component.
Tetanus (Lockjaw) – An acute, often fatal disease caused by bacterial toxins. It causes painful tightening of the muscles, usually all over the body. It can lead to “locking” of the jaw, stiff neck, difficulty swallowing, and stiff, painful muscle contractions.	Approximately 100% protection against the disease shortly after vaccination. However, immunity decreases with time, so it is important to receive booster vaccinations.	Same as for DTaP/Tdap vaccine (see diphtheria).
Pertussis (Whooping Cough) – A bacteria which causes bursts of rapid cough followed by a long inspiratory effort accompanied by a characteristic high-pitched whoop. During such an attack, the infected person often turns blue from a lack of oxygen, they also may look very ill or distressed. This can be followed by vomiting and exhaustion. These spells can last for weeks or months.	Estimates of vaccine efficacy are around 80% shortly after vaccination. However, immunity decreases over time, so it is important to receive booster vaccinations.	Same as for DTaP/Tdap vaccine (see diphtheria).
Polio – A virus which enters the body through the mouth. Sometimes, it does not cause serious illness, but it can cause paralysis (can't move arms or legs); muscle function may or may not return. It can cause death by paralyzing the muscles that allow breathing.	More than 90% of vaccine recipients become immune with just two doses. After receiving three doses, 99% will become immune to the disease.	IPV vaccine – Minor local reactions such as pain and redness at the injection site may occur, but are uncommon. No serious reactions to the vaccine have been reported.
Measles – A virus causes an infection that primarily occurs in the respiratory system, but later causes rash, cough, runny nose, eye redness, and fever. It can lead to ear infections, pneumonia, seizures (jerking and staring), brain damage, and death.	Measles protection develops in approximately 95% of children vaccinated at 12 months of age and 98% of children vaccinated at 15 months of age. Two doses of vaccine provide protection in more than 99% of those vaccinated.	MMR vaccine – The most common side effects of vaccination in children are fever and rash. If these problems occur, it is usually within seven to 12 days after vaccination and lasts one or two days. There is no evidence linking the MMR vaccine to autism.

Vaccine-preventable disease	Effectiveness of vaccine	Possible vaccine side effects
Mumps – A virus which causes a fever, headache, swollen lymph nodes (especially on the face and neck), muscle pain, and fatigue. It can lead to deafness, meningitis (inflammation surrounding the brain and spinal cord), painful swelling of the testicles or ovaries leading to sterility (inability to have children), and rarely death.	Two doses of MMR is estimated to be 88% effective against mumps.	Same as for MMR vaccine (see Measles).
Rubella – A virus which causes a rash, mild fever, fatigue, and arthritis (joint swelling). If a woman gets rubella while pregnant, she could have a miscarriage or a stillbirth. The baby may also be born with serious birth defects.	More than 95% of those vaccinated at 12 months of age and older developed protection against the virus. Protection will last for at least 15 years in more than 90% of vaccinated persons.	Same as for MMR vaccine (see Measles).
Hepatitis B – An acute liver infection caused by a virus which can provoke long-term (chronic) illness that leads to liver damage (cirrhosis), liver cancer, and death.	The vaccine is 80 to 100% effective in preventing infection or clinical hepatitis in those who receive the full series of the vaccine.	The most common side effects include pain at the injection site, fatigue, and headache. Fevers are uncommon, while severe reactions are very rare.
Hepatitis A – An acute liver infection caused by a virus which can lead to symptoms ranging from mild to severe, including “flu-like” illness, loss of appetite, nausea, diarrhea, severe stomach pains, and jaundice (yellow skin or eyes). In rare cases, hepatitis A can cause fulminant hepatitis (acute liver failure) or death.	After the first dose, 97% of children and adolescents will develop protection within four weeks. After receiving the second dose, 100% will be protected against the virus.	The most common side effects are pain, redness, and swelling at the injection site. Fatigue and low-grade fever are reported in fewer than 10% of persons, lasting no more than one or two days. Serious allergic reactions are very rare.
Varicella (Chickenpox) – A virus which causes generalized rash, itching, fever, and fatigue. It can lead to severe skin infections, scars, pneumonia, brain damage, or death. Varicella can return later in life as Shingles.	After one dose of vaccine, 97 to 99% of children develop protection against the virus. More than 90% of vaccine responders maintain protection for at least seven to 10 years. Vaccine efficacy is estimated to be 70 to 90% against infection, and 90 to 100% against severe disease.	The most common side effects are pain and redness at the injection site, and mild rash. Reactions may be more severe in immunocompromised people.
Haemophilus influenzae type b (Hib) – A bacteria enters the body and causes invasive diseases such as meningitis (inflammation surrounding the brain and spinal cord), pneumonia, severe swelling in the throat (making it hard to breathe), arthritis, and death.	More than 95% of infants will develop protection against these bacteria after a series of two or three doses. The efficacy of the vaccine in preventing clinical cases of the disease is estimated to be 95 to 100%. Children with chronic diseases such as sickle cell disease and HIV infection seem to respond well to Hib vaccines.	The most common side effects after vaccination are swelling, redness, or pain at the injection site, resolving in less than 24 hours. About one in 20 people have discomfort or local inflammation. Fever and irritability are not frequent side effects. Serious reactions are uncommon.

Vaccine-preventable disease	Effectiveness of vaccine	Possible vaccine side effects
<p>Meningococcal disease – A bacteria that causes an acute, potentially severe illness that most often leads to meningitis, an infection of the spinal fluid and the fluid that surrounds the brain. These bacteria have at least 13 different subtypes (serogroups). Five of these serogroups, A, B, C, Y, and W, cause almost all invasive disease. It leads to sudden onset of fever, headache, and stiff neck and is usually accompanied by nausea, vomiting, light sensitivity, and altered mental status.</p> <p>Less commonly, it can cause pneumonia, blood infection, arthritis, and ear / throat infections. Meningococcal disease can result in hearing loss, nervous system problems, seizures, strokes, loss of limbs (arms, legs), and death.</p>	<p>MenACWY vaccine protects against four types of meningococcal disease. MenACWY vaccine effectiveness may wane over time. Vaccine effectiveness is estimated at 79% among adolescents vaccinated within the past year and is estimated at 61% effectiveness three to eight years after vaccination. Children with chronic diseases such as sickle cell disease and HIV infection seem to respond well to MenACWY vaccines.</p> <p>Immune response for MenB vaccine suggests the vaccines are protective, but effectiveness data is limited. Since meningococcal disease is uncommon, many people need to get these vaccines in order to measure their effectiveness. Since the CDC first began recommending Meningococcal vaccines in 2005, incidence of the disease has decreased by more than 90%.</p>	<p>The most common side effects of MenACWY and MenB vaccines are redness or pain at the injection site lasting between one and two days, headache, and fatigue. Serious allergic reactions are very rare.</p>
<p>Pneumococcal disease – A bacteria that causes an infection which can lead to blood infections, pneumonia, and bacterial meningitis (swelling of the lining of the brain). Pneumococcal pneumonia is the most common infection, leading to abrupt onset of fever, chills, chest pain, cough, and breathing difficulty. Complications include swelling of the sac around the heart and infection of the area around the lungs. Pneumococcal meningitis kills about 14% of people who develop it. Pneumococcal meningitis can also lead to other health problems, including deafness and brain damage.</p>	<p>Vaccination has been shown to be 97% effective in preventing invasive disease caused by bacteria contained in the vaccine and 89% effective against all types of <i>S. pneumoniae</i>, including those not found in the vaccine. Children with chronic diseases such as sickle cell disease and HIV infection also seem to respond well to Pneumococcal conjugate vaccine (PCV15, PCV20). Pneumococcal polysaccharide vaccine (PPSV23) (<i>recommended for individuals with certain high risk factors and adults ≥ 65 years</i>) is 60 to 70% effective in preventing invasive disease.</p>	<p>The most common side effects are local reactions such as redness, tenderness, or swelling at the injection site. Other side effects such as fever and muscle pain are uncommon and severe reactions are rare.</p>

Note: The Rabies vaccine isn't required for K-12 education but is necessary for veterinary school enrollment.

The Texas Department of State Health Services (DSHS) Immunization Exemption for Reason of Conscience Affidavit Form is utilized by various entities, including Texas child care facilities, schools and higher education institutions.

Texas Department of State Health Services
Immunization Section



Stock Number 11-11846
Rev. 07/2024

Beneficios y riesgos de la vacunación

Enfermedades prevenibles con vacuna	Eficacia de la vacuna	Posibles efectos secundarios de vacunación
Difteria – Las toxinas de las bacterias causan la formación de una gruesa capa en las membranas mucosas, con mayor frecuencia en la parte posterior de la garganta. Puede ocasionar problemas de respiración, parálisis, insuficiencia cardiaca, coma y la muerte en el curso de seis a 10 días.	Se calcula que el toxoide diftérico tiene una eficacia clínica del 97% contra la enfermedad.	Vacuna DTaP – Los efectos secundarios más comunes incluyen enrojecimiento, dolor e hinchazón en el sitio de la inyección. La presencia de fiebre no es común, y ocurre en menos del 5% del tiempo. Las reacciones graves son raras.
Tétanos – Enfermedad aguda, a menudo mortal, causada por la toxina de la bacteria del tétanos. Causa rigidez dolorosa de los músculos, normalmente en todo el cuerpo. Puede occasionar el “cierre” de la mandíbula, fuerte rigidez del cuello, dificultad para tragar e intensas, dolorosas contracciones musculares.	Protección de aproximadamente 100% contra la enfermedad.	Los mismos efectos secundarios que para la vacuna DTaP (véase Difteria).
Tos ferina (Pertussis) – Causa rápidos accesos de tos que terminan con dificultad para respirar. Las personas infectadas a menudo adquieren una coloración azul debido a la falta de aire. Dichos accesos pueden durar de semanas a meses. Puede occasionar neumonía, ataques convulsivos (episodios de sacudidas musculares y de tener la mirada perdida), daño cerebral e incluso la muerte.	Se calcula que la eficacia de la vacuna es del 80 al 85% para las vacunas actualmente autorizadas en Estados Unidos.	Los mismos efectos secundarios que para la vacuna DTaP (véase Difteria).
Polio – Se introduce en el cuerpo a través de la boca. A veces no causa una enfermedad grave, pero puede causar parálisis (no poder mover los brazos o las piernas); la función muscular podría o no restituirse. Puede causar la muerte al paralizar los músculos que intervienen en la respiración.	Más de 90% de quienes reciben la vacuna se vuelven inmunes con solo dos dosis. Después de haber recibido tres dosis, el 99% se volverán inmunes a la enfermedad.	Vacuna IPV – Pueden ocurrir reacciones localizadas menores, como dolor y enrojecimiento en el sitio de la inyección, pero no son comunes. No se ha informado de reacciones graves a la vacuna.
Sarampión – La infección ocurre principalmente en el aparato respiratorio, pero posteriormente causa sarpullido, tos, moco nasal, irritación de los ojos y fiebre. Puede occasionar infecciones de los oídos, neumonía, ataques convulsivos (con sacudidas musculares y la mirada perdida), daño cerebral y la muerte.	La protección contra el sarampión se desarrolla en aproximadamente 95% de los niños vacunados a los 12 meses, y en el 98% de los niños vacunados a los 15 meses. Dos dosis de la vacuna proveen protección en más del 99% .	Vacuna MMR – Los efectos secundarios más comunes de la vacuna en los niños son la fiebre y el sarpullido. De ocurrir, estos problemas aparecen normalmente entre siete y 12 días después de la vacunación y duran de uno o dos días. No existe evidencia que vincule la vacuna MMR con el autismo.

Enfermedades prevenibles con vacuna	Eficacia de la vacuna	Posibles efectos secundarios de vacunación
Paperas – Causan fiebre, dolor de cabeza, inflamación de los ganglios, dolor muscular y cansancio. Pueden ocasionar sordera, meningitis (inflamación alrededor del cerebro y la médula espinal) e inflamación dolorosa de testículos u ovarios, la cual ocasiona esterilidad (incapacidad para tener hijos); rara vez ocasionan la muerte.	Se calcula que dos dosis de la vacuna MMR tienen una eficacia del 88% contra las paperas.	Los mismos efectos secundarios que para la vacuna MMR (véase Sarampión).
Rubeola – Causa sarpullido, fiebre ligera, cansancio y artritis (inflamación de las articulaciones). Si a una mujer le da rubeola estando embarazada, podría tener un aborto espontáneo, o su bebé podría nacer muerto o tener graves defectos de nacimiento.	Más del 95% de los vacunados a los 12 meses de edad o mayores desarrollaron protección contra el virus. La protección dura al menos 15 años en más del 90% de las personas vacunadas.	Los mismos efectos secundarios que para la vacuna MMR (véase Sarampión).
Hepatitis B – Afección del hígado que causa enfermedad a largo plazo (crónica) y da lugar a daño al hígado (cirrosis), cáncer del hígado e incluso la muerte.	La vacuna tiene una eficacia del 80% al 100% en la prevención de la infección o la hepatitis clínica en aquellos que reciben la serie completa de la vacuna.	Los efectos secundarios más comunes incluyen dolor en el sitio de la inyección, cansancio y dolor de cabeza. La fiebre no es común, y las reacciones graves son muy raras.
Hepatitis A – Afección del hígado que causa una enfermedad de leve a grave parecida a la gripe, ictericia (amarillez de la piel o los ojos), intenso dolor de estómago, pérdida del apetito, náusea y diarrea.	Después de la primera dosis, el 97% de personas desarrollan protección en el curso de cuatro semanas. Después de recibir la segunda dosis, el 100% están protegidos contra el virus.	Los efectos secundarios más comunes son el dolor, el enrojecimiento y la hinchazón en el sitio de la inyección. En menos del 10% de las personas se informa de cansancio y fiebre moderada que no duran más de uno o dos días. Las reacciones alérgicas graves son muy raras.
Varicela – Infección aguda que causa un sarpullido generalizado, comezón, fiebre y cansancio. Puede ocasionar graves infecciones de la piel, cicatrices, neumonía, daño cerebral o la muerte. Una vez infectada la persona, la varicela se puede reactivar en forma de herpes zóster más adelante con el tiempo.	Tras una dosis de la vacuna, entre el 97 y 99% de los niños desarrollan protección contra el virus. Más del 90% de aquellos que responden a la vacuna conservan la protección por al menos Siete a 10 años. Se calcula que la eficacia de la vacuna es del 70 al 90% contra la infección, y del 90 al 100% contra la enfermedad grave.	Los efectos secundarios más comunes son el dolor y el enrojecimiento en el sitio de la inyección, y un sarpullido leve. Las reacciones podrían ser más graves en las personas inmunodeprimidas.

Enfermedades prevenibles con vacuna	Eficacia de la vacuna	Posibles efectos secundarios de vacunación
<i>Haemophilus influenzae tipo b (Hib)</i> – Causa enfermedades invasivas como la meningitis (inflamación alrededor del cerebro y la médula espinal), neumonía, grave inflamación de la garganta (dificultando la respiración), artritis y la muerte.	Más del 95% de los bebés desarrollan protección contra estas bacterias después de una serie de dos o tres dosis. Se calcula que la eficacia de la vacuna en la prevención de casos clínicos de la enfermedad es del 95 al 100%.	Los efectos secundarios más comunes después de la vacunación son la inflamación, el enrojecimiento o el dolor en el sitio de la inyección, los cuales duran menos de 24 horas. Alrededor de 1 de cada 20 personas tiene malestar o inflamación localizada. La fiebre y la irritabilidad no son efectos secundarios frecuentes. Las reacciones graves no son comunes.
Enfermedad meningocócica – Es una enfermedad aguda, potencialmente grave, que con gran frecuencia causa meningitis, una infección del líquido cefalorraquídeo y el líquido que envuelve el cerebro. Causa una rápida aparición de fiebre, dolor de cabeza e intensa rigidez del cuello, y generalmente se acompaña de náusea, vómito, sensibilidad a la luz y alteración del estado mental. Con menor frecuencia, puede causar neumonía, infección de la sangre, artritis e infecciones del oído o la garganta. La enfermedad meningocócica puede ocasionar pérdida del oído, problemas del sistema nervioso, ataques convulsivos, derrame cerebral, pérdida de extremidades (brazos, piernas) y la muerte.	La vacuna MenACWY protege contra cuatro tipos de enfermedad meningocócica. La eficacia de la vacuna MenACWY puede disminuir con el tiempo. Se calcula que la eficacia de la vacuna es del 82% entre los adolescentes que se vacunaron en el último año, y se calcula que es del 69% para los que se vacunaron dentro de los seis últimos años.	Los efectos secundarios más comunes de la vacuna MenACWY son el enrojecimiento o el dolor en el sitio de la inyección durante uno o dos días, dolor de cabeza y cansancio. Las reacciones alérgicas graves son muy raras.
Enfermedad neumocócica – Infección por bacterias que puede causar infecciones en la sangre, neumonía y meningitis bacteriana (inflamación del revestimiento del cerebro). La neumonía neumocócica es la infección más común, la cual causa una aparición repentina de fiebre, escalofríos, dolor en el pecho, tos y dificultad para respirar. Las complicaciones incluyen la inflamación del saco que rodea al corazón e infección del área alrededor de los pulmones. La meningitis neumocócica causa la muerte aproximadamente a 30% de las personas que la contraen. La meningitis neumocócica también puede ocasionar otros problemas de salud, incluida la sordera y el daño cerebral.	La vacunación ha demostrado una eficacia del 97% en la prevención de la enfermedad invasiva causada por la bacteria contenida en la vacuna, y una eficacia del 89% contra todos los tipos de <i>S. pneumoniae</i> , incluidos aquellos que no se encuentran en la vacuna. Los niños con enfermedades crónicas como la enfermedad de células falciformes y la infección por VIH también parecen responder bien a la vacuna antineumocócica conjugada (PCV13). La vacuna antineumocócica polisacárida (PPSV) (recomendada para individuos con ciertos factores de alto riesgo y adultos de 65 años de edad o mayores) tiene una efectividad del 60 al 70% en la prevención de la enfermedad invasiva.	Los efectos secundarios más comunes son las reacciones localizadas como enrojecimiento, sensibilidad o inflamación en el sitio de la inyección. Otros efectos secundarios como la fiebre y el dolor muscular no son comunes y las reacciones graves son raras.

Texas Department of State Health Services
Immunization Section



Stock Number 11-11846
Rev. 07/2024