

LOW INFORMATION,
LOW AWARENESS

fatalism
NEGATIVE HEALTH BELIEFS

LOW LEVELS OF SOCIAL SUPPORT



HIGH LEVELS OF STIGMA

“I don’t know where to start,
LOW SELF-EFFICACY
all of this is too overwhelming”

LOW LEVELS OF ENGAGEMENT
WITH HEALTH CARE SYSTEMS



LOW EDUCATIONAL ATTAINMENT

“INVINCIBLES”

“I feel healthy”
A red hand holding a red heart.

<25 YOUNGER PEOPLE
THINK LESS ABOUT
LONG TERM HEALTH

“I don’t have AIDS so
I don’t need to see a
doctor until then”
AIDS”

HIGH CD4 COUNT
NO AIDS DIAGNOSIS

Not feeling sick is
the number 1 rea-
son PLWH give for
not accessing care

Dropping
OUT OF
CARE risk
factors

LACK OF STEADY EMPLOYMENT/
HOUSING/TRANSPORTATION



LOW LEVELS OF SOCIAL SUPPORT /
HIGH STIGMA




One out of five Black gay men
will drop out of care for 12 or
more months

LOW INCOME



LACK OF
HEALTH INSURANCE

MINORITY RACE/ETHNICITY



56%

of women out
of care are
Black Women

VULNERABLE POPULATIONS

INJECTION DRUG USE



UNTREATED
MENTAL
HEALTH
ISSUES



CHRONIC MISSED
APPOINTMENTS

HEPATITIS C

OTHER COMORBIDITIES

COMPETING PRIORITIES