Collaborative Staging Codes
Glottic Larynx
C32.0
C32.0 Glottis

Glottic Larynx
CS Tumor Size
SEE STANDARD TABLE

Glottic Larynx
CS Extension (Revised: 08/14/2006)

Code	Description	TNM	SS77	SS2000
00	In situ; noninvasive; intraepithelial	Tis	IS	IS
10	Invasive tumor with normal vocal cord mobility Confined to glottis, NOS; intrinsic larynx; laryngeal commisure(s) anterior, posterior; vocal cord(s), NOS, true vocal cord(s), true cord(s)	T1NOS	L	L
11	One vocal cord	T1a	L	L
12	Both vocal cords	T1b	L	L
30	Tumor involves adjacent regions(s) of larynx Subglottis Supraglottis False vocal cord (s)	T2	L	L
35	Impaired vocal cord mobility	T2	L	L
40	Tumor limited to larynx WITH vocal cord fixation Involvement of intrinsic muscle(s): Aryepiglottic Corniculate tubercle Cuneiform tubercle Arytenoid Cricoarytenoid Cricothyroid Thyroarytenoid Thyroepiglottic Vocalis	T3	L	L
45	Localized, NOS	T1NOS	L	L
51	Paraglottic space	T3	RE	RE

52	Minor thyroid cartilage erosion (e.g., inner cortex)	T3	RE	D
60	Base of tongue Hypopharynx, NOS Pre-epiglottic tissues Postcricoid area Pyriform sinus Vallecula	T4a	RE	RE
68	Extension to/through Cricoid cartilage Thyroid cartilage except minor erosion, see code 52	T4a	RE	D
70	Extension to/through tissues beyond larynx: Extrinsic (strap) muscles Omohyoid Sternohyoid Sternothyroid Thryohyoid Oropharynx Skin Soft tissue of neck Thyroid gland Trachea	T4a	D	D
71	Cervical esophagus	T4a	D	D
73	Deep extrinsic muscle(s) of tongue	T4a	D	D
80	Further contiguous extension, including: Mediastinal structures Prevertebral space Tumor encases carotid artery	T4b	D	D
95	No evidence of primary tumor	T0	U	U
99	Unknown extension Primary tumor cannot be assessed Not documented in patient record	TX	U	U

Glottic Larynx
CS TS/Ext-Eval
SEE STANDARD TABLE
NOTE: For this primary site code CS Tumor Size/Ext Eval on the basis of the CS Extension field only.

Glottic Larynx
CS Lymph Nodes (Revised: 08/21/2006)
Note 1: For head and neck schemas, this field includes all lymph nodes defined as Levels I-VII and Other by AJCC.
Note 2: For head and neck schemas, additional information about lymph nodes (size of involved nodes, extracapsular extension, and levels involved) is coded in Site-Specific Factors 1-6.
Note 3: If laterality of lymph nodes is not specified, assume nodes are ipsilateral. Midline nodes are considered ipsilateral.
Note 4: For head and neck cancers, if lymph nodes are described only as "supraclavicular", try to determine if they are in Level IV (deep to the sternocleidomastoid muscle, in the lower jugular chain) or Level V (in the posterior triangle, inferior to the transverse cervical artery) and code appropriately. If the specific level cannot be determined, consider them as Level V nodes.

Code	Description	TNM	SS77	SS2000
00	None; no regional lymph node involvement	N0	NONE	NONE
10	Single positive ipsilateral regional node: Level II Jugulodigastric (subdigastric) Upper deep cervical Upper jugular Level III Middle deep cervical Mid-jugular Level IV Jugulo-omohyoid (supraomohyoid) Lower deep cervical Lower jugular Level VI Anterior deep cervical Delphian node Laterotracheal Paralaryngeal Paratracheal Prelaryngeal (Delphian) Pretracheal Recurrent laryngeal Cervical, NOS Deep cervical, NOS Internal jugular NOS: Regional lymph node, NOS Stated as N1, NOS	*	RN	RN

11	Single positive ipsilateral regional node: Level I Submandibular (submaxillary) Submental Other groups Retropharyngeal Mandibular, NOS	*	D	RN
12	Single positive ipsilateral regional node: Level V node Posterior cervical Posterior triangle (spinal accessory and transverse cervical) (upper, middle, and lower corresponding to the levels that define upper, middle, and lower jugular nodes) Level VII node Upper mediastinum (for other mediastinal nodes see CS Mets at DX) Other groups Intraparotid Parapharyngeal Periparotid Sub-occipital Supraclavicular, NOS (See Note 4)	*	D	RN
18	Stated as N1, no other information	N1	RN	RN
19	Stated as N2a, no other information	N2a	RN	RN
20	Multiple positive ipsilateral nodes listed in code 10	*	RN	RN
21	Multiple positive ipsilateral nodes listed in code 11	*	D	RN
22	Multiple positive ipsilateral nodes listed in code 12	*	D	D
29	Stated as N2b, no other information	N2b	RN	RN
30	Regional lymph nodes as listed in code 10: Positive ipsilateral node(s), not stated if single or multiple	*	RN	RN
31	Regional lymph nodes as listed in code 11: Positive ipsilateral node(s), not stated if single or multiple	*	D	RN
32	Regional lymph nodes as listed in code 12: Positive ipsilateral node(s), not stated if single or multiple	*	D	D

40	Regional lymph nodes as listed in code 10: Positive bilateral or contralateral nodes	*	RN	RN
41	Regional lymph nodes as listed in code 11: Positive bilateral or contralateral nodes	*	D	RN
42	Regional lymph nodes as listed in code 12: Positive bilateral or contralateral nodes	*	D	D
49	Stated as N2c, no other information	N2c	RN	RN
50	Regional lymph nodes as listed in code 10: Positive node(s) not stated if ipsilateral, or bilateral, or contralateral AND not stated if single or multiple	*	RN	RN
51	Regional lymph nodes as listed in code 11: Positive node(s) not stated if ipsilateral, or bilateral, or contralateral AND not stated if single or multiple	*	D	RN
52	Regional lymph nodes as listed in code 12: Positive node(s), not stated if ipsilateral, or bilateral, or contralateral AND not stated if single or multiple	*	D	D
60	Stated as N2NOS	N2NOS	RN	RN
70	Stated as N3, no other information	N3	RN	RN
80	Lymph nodes, NOS, no other information	*	RN	RN
99	Unknown; not stated Regional lymph node(s) cannot be assessed Not documented in patient record	NX	U	U

Glottic Larynx
Reg LN Pos
SEE STANDARD TABLE

Glottic Larynx
Reg LN Exam
SEE STANDARD TABLE

Glottic Larynx
CS Mets at DX (Revised: 08/14/2006)
Note: Supraclavicular and transverse cervical lymph nodes are now coded in CS Lymph Nodes because they are categorized as N rather than M in AJCC TNM. Any cases coded to 10 or 50 can be reviewed and recoded. The volume of cases affected should be small.

Code	Description	TNM	SS77	SS2000
00	No; none	M0	NONE	NONE
10	Distant lymph node(s) Mediastinal Supraclavicular (transverse cervical) Distant lymph node(s), NOS	M1	D	D
40	Distant metastases except distant lymph node(s) (code 10) Distant metastasis, NOS Carcinomatosis	M1	D	D
50	(10) + (40) Distant lymph node(s) plus other distant metastases	M1	D	D
99	Unknown if distant metastasis Distant metastasis cannot be assessed Not documented in patient record	MX	U	U

Collaborative Staging Codes
Supraglottic Larynx
C32.1
C32.1 Supraglottis
Note: Excludes Anterior Surface of Epiglottis - see separate schema (C10.1).

Supraglottic Larynx
CS Tumor Size
SEE STANDARD TABLE

Supraglottic Larynx
CS Extension (Revised: 03/17/2004)

Code	Description	TNM	SS77	SS2000
00	In situ; noninvasive; intraepithelial	Tis	IS	IS
10	Invasive tumor with normal vocal cord mobility confined to: Supraglottis (one subsite): Aryepiglottic fold Arytenoid cartilage Corniculate cartilage Cuneiform cartilage Epilarynx, NOS False cords Ventricular bands Ventricular cavity Ventricular fold Infrahyoid epiglottis Laryngeal cartilage, NOS Laryngeal (posterior) surface of epiglottis Suprahyoid epiglottis (including tip, lingual {anterior} and laryngeal surfaces)	T1	L	L
20	Tumor involves more than one subsite of supraglottis WITHOUT fixation or NOS	T2	L	L
30	Tumor involves adjacent regions(s) of larynx	T2	L	L
35	Impaired vocal cord mobility	T2	L	L
40	Tumor limited to larynx WITH vocal cord fixation	T3	L	L
45	Localized, NOS	T1	L	L
52	Paraglottic space	T3	RE	RE

60	Tumor involves region outside the supraglottis WITHOUT fixation, including: Medial wall of pyriform sinus Mucosa of base of tongue Vallecula	T2	RE	RE
62	Code 60 WITH fixation	T3	RE	RE
65	Hypopharynx, NOS Postcricoid area Pre-epiglottic tissues	T3	RE	RE
66	Deep base of tongue	T3	RE	RE
67	Cricoid cartilage	T3	RE	RE
68	Minor thyroid cartilage erosion (e.g., inner cortex)	T3	RE	D
70	Extension to/through: Esophagus Oropharynx Soft tissues of neck Thyroid cartilage (except minor erosion, see code 68) Thyroid gland	T4a	D	D
72	Extension to/through: Extrinsic (strap) muscle(s) Omohyoid Sternohyoid Sternothyroid Thyrohyoid Skin	T4a	D	D
73	Extension to/through: Deep extrinsic muscle of tongue Trachea	T4a	D	D
80	Further contiguous extension, including: Mediastinal structures Prevertebral space Tumor encases carotid artery	T4b	D	D
95	No evidence of primary tumor	T0	U	U
99	Unknown extension Primary tumor cannot be assessed Not documented in patient record	TX	U	U

Supraglottic Larynx
CS TS/Ext-Eval
SEE STANDARD TABLE
NOTE: For this primary site code CS Tumor Size/Ext Eval on the basis of the CS Extension field only.

Supraglottic Larynx
CS Lymph Nodes (Revised: 08/21/2006)
Note 1: For head and neck schemas, this field includes all lymph nodes defined as Levels I-VII and Other by AJCC.
Note 2: For head and neck schemas, additional information about lymph nodes (size of involved nodes, extracapsular extension, and levels involved) is coded in Site-Specific Factors 1-6.
Note 3: If laterality of lymph nodes is not specified, assume nodes are ipsilateral. Midline nodes are considered ipsilateral.
Note 4: For head and neck cancers, if lymph nodes are described only as "supraclavicular", try to determine if they are in Level IV (deep to the sternocleidomastoid muscle, in the lower jugular chain) or Level V (in the posterior triangle, inferior to the transverse cervical artery) and code appropriately. If the specific level cannot be determined, consider them as Level V nodes.

Code	Description	TNM	SS77	SS2000
00	None; no regional lymph node involvement	N0	NONE	NONE
10	Single positive ipsilateral regional node: Level II Jugulodigastric (subdigastric) Upper deep cervical Upper jugular Level III Middle deep cervical Mid-jugular Level VI Anterior deep cervical Delphian node Laterotracheal Paralaryngeal Paratracheal Prelaryngeal Pretracheal Recurrent laryngeal Cervical, NOS Deep cervical, NOS Internal jugular, NOS: Regional lymph node, NOS Stated as N1, NOS	*	RN	RN

11	Single positive ipsilateral regional node: Level I Submandibular (submaxillary) Submental Other groups Retropharyngeal Mandibular, NOS	*	D	RN
12	Single positive ipsilateral regional node: Level V node Posterior cervical Posterior triangle (spinal accessory and transverse cervical) (upper, middle, and lower corresponding to the levels that define upper, middle, and lower jugular nodes) Level VII node Upper mediastinum (for other mediastinal nodes see CS Mets at DX) Other groups Intraparotid Parapharyngeal Periparotid Sub-occipital Supraclavicular, NOS (See note 4)	*	D	D
18	Stated as N1, no other information	N1	RN	RN
19	Stated as N2a, no other information	N2a	RN	RN
20	Multiple positive ipsilateral nodes listed in code 10	*	RN	RN
21	Multiple positive ipsilateral nodes listed in code 11	*	D	RN
22	Multiple positive ipsilateral nodes listed in code 12	*	D	D
29	Stated as N2b, no other information	N2b	RN	RN
30	Regional lymph nodes as listed in code 10: Positive ipsilateral node(s), not stated if single or multiple	*	RN	RN
31	Regional lymph nodes as listed in code 11: Positive ipsilateral node(s), not stated if single or multiple	*	D	RN
32	Regional lymph nodes as listed in code 12: Positive ipsilateral node(s), not stated if single or multiple	*	D	D

40	Regional lymph nodes as listed in code 10: Positive bilateral or contralateral nodes	*	RN	RN
41	Regional lymph nodes as listed in code 11: Positive bilateral or contralateral nodes	*	D	RN
42	Regional lymph nodes as listed in code 12: Positive bilateral or contralateral nodes	*	D	D
49	Stated as N2c, no other information	N2c	RN	RN
50	Regional lymph nodes as listed in code 10: Positive node(s) not stated if ipsilateral, or bilateral, or contralateral AND not stated if single or multiple	*	RN	RN
51	Regional lymph nodes as listed in code 11: Positive node(s) not stated if ipsilateral, or bilateral, or contralateral AND not stated if single or multiple	*	D	RN
52	Regional lymph nodes as listed in code 12: Positive node(s), not stated if ipsilateral, or bilateral, or contralateral AND not stated if single or multiple	*	D	D
60	Stated as N2, NOS	N2NOS	RN	RN
70	Stated as N3, no other information	N3	RN	RN
80	Lymph nodes, NOS, no other information	*	RN	RN
99	Unknown; not stated Regional lymph node(s) cannot be assessed Not documented in patient record	NX	U	U

Supraglottic Larynx
Reg LN Pos
SEE STANDARD TABLE

Supraglottic Larynx
Reg LN Exam
SEE STANDARD TABLE

Supraglottic Larynx
CS Mets at DX (Revised: 08/14/2006)
Note: Supraclavicular and transverse cervical lymph nodes are now coded in CS Lymph Nodes because they are categorized as N rather than M in AJCC TNM. Any cases coded to 10 or 50 can be reviewed and recoded. The volume of cases affected should be small.

Code	Description	TNM	SS77	SS2000
00	No; none	M0	NONE	NONE
10	Distant lymph node(s) Mediastinal Distant lymph node(s), NOS	M1	D	D
40	Distant metastases except distant lymph node(s) (code 10) Distant metastasis, NOS Carcinomatosis	M1	D	D
50	(10) + (40) Distant lymph node(s) plus other distant metastases	M1	D	D
99	Unknown if distant metastasis Distant metastasis cannot be assessed Not documented in patient record	MX	U	U

Collaborative Staging Codes
Subglottic Larynx
C32.2
C32.2 Subglottis

Subglottic Larynx
CS Tumor Size
SEE STANDARD TABLE

Subglottic Larynx
CS Extension (Revised: 03/17/2004)

Code	Description	TNM	SS77	SS2000
00	In situ; noninvasive; intraepithelial	Tis	IS	IS
10	Invasive tumor with normal vocal cord mobility confined to subglottis	T1	L	L
30	Tumor involves adjacent regions(s) of larynx Vocal cords with normal or impaired mobility	T2	L	L
40	Tumor limited to larynx WITH vocal cord fixation	T3	L	L
45	Localized, NOS	T1	L	L
60	Base of tongue Hypopharynx, NOS Postcricoid area Pre-epiglottic tissues Pyriform sinus (pyriform fossa) Vallecula	T4a	RE	RE
68	Extension to/through cricoid cartilage or thyroid cartilage	T4a	RE	D
70	Extension to/through: Cervical esophagus Deep extrinsic muscles of tongue Extrinsic (strap) muscles Omohyoid Sternohyoid Sternothyroid Thyrohyoid Oropharynx Skin Soft tissues of neck Thyroid gland Trachea	T4a	D	D

73	Contiguous extension to other tissues beyond larynx not specified in codes 70 or 80	T4a	D	D
80	Further contiguous extension: Mediastinal structures Prevertebral space Tumor encases carotid artery	T4b	D	D
95	No evidence of primary tumor	T0	U	U
99	Unknown extension Primary tumor cannot be assessed Not documented in patient record	TX	U	U

Subglottic Larynx
CS TS/Ext-Eval
SEE STANDARD TABLE
NOTE: For this primary site code CS Tumor Size/Ext Eval on the basis of the CS Extension field only.

Subglottic Larynx
CS Lymph Nodes (Revised: 08/21/2006)
Note 1: For head and neck schemas, this field includes all lymph nodes defined as Levels I-VII and Other by AJCC.
Note 2: For head and neck schemas, additional information about lymph nodes (size of involved nodes, extracapsular extension, and levels involved) is coded in Site-Specific Factors 1-6.
Note 3: If laterality of lymph nodes is not specified, assume nodes are ipsilateral. Midline nodes are considered ipsilateral.
Note 4: For head and neck cancers, if lymph nodes are described only as "supraclavicular", try to determine if they are in Level IV (deep to the sternocleidomastoid muscle, in the lower jugular chain) or Level V (in the posterior triangle, inferior to the transverse cervical artery) and code appropriately. If the specific level cannot be determined, consider them as Level V nodes.

Code	Description	TNM	SS77	SS2000
00	None; no regional lymph node involvement	N0	NONE	NONE
10	Single positive ipsilateral regional node: Level II Jugulodigastric (subdigastric) Upper deep cervical Upper jugular Level III Middle deep cervical Mid-jugular Level VI Anterior deep cervical	*	RN	RN

10 cont'd	Delphian node Laterotracheal Paralaryngeal Paratracheal Prelaryngeal Pretracheal Recurrent laryngeal Cervical, NOS Deep cervical, NOS Internal jugular, NOS: Regional lymph node, NOS Stated as N1, NOS	*	RN	RN
11	Single positive ipsilateral regional node: Level I Submandibular (submaxillary) Submental Other groups Retropharyngeal Mandibular, NOS	*	D	RN
12	Single positive ipsilateral regional node: Level V node Posterior cervical Posterior triangle (spinal accessory and transverse cervical) (upper, middle, and lower corresponding to the levels that define upper, middle, and lower jugular nodes) Level VII node Upper mediastinum (for other mediastinal nodes see CS Mets at DX) Other groups Intraparotid Parapharyngeal Periparotid Sub-occipital Supraclavicular, NOS (See note 4)	*	D	D
18	Stated as N1, no other information	N1	RN	RN
19	Stated as N2a, no other information	N2a	RN	RN
20	Multiple positive ipsilateral nodes listed in code 10	*	RN	RN
21	Multiple positive ipsilateral nodes listed in code 11	*	D	RN
22	Multiple positive ipsilateral nodes listed in code 12	*	D	D

29	Stated as N2b, no other information	N2b	RN	RN
30	Regional lymph nodes as listed in code 10: Positive ipsilateral node(s), not stated if single or multiple	*	RN	RN
31	Regional lymph nodes as listed in code 11: Positive ipsilateral node(s), not stated if single or multiple	*	D	RN
32	Regional lymph nodes as listed in code 12: Positive ipsilateral node(s), not stated if single or multiple	*	D	D
40	Regional lymph nodes as listed in code 10: Positive bilateral or contralateral nodes	*	RN	RN
41	Regional lymph nodes as listed in code 11: Positive bilateral or contralateral nodes	*	D	RN
42	Regional lymph nodes as listed in code 12: Positive bilateral or contralateral nodes	*	D	D
49	Stated as N2c, no other information	N2c	RN	RN
50	Regional lymph nodes as listed in code 10: Positive node(s) not stated if ipsilateral, or bilateral, or contralateral AND not stated if single or multiple	*	RN	RN
51	Regional lymph nodes as listed in code 11: Positive node(s) not stated if ipsilateral, or bilateral, or contralateral AND not stated if single or multiple	*	D	RN
52	Regional lymph nodes as listed in code 12: Positive node(s), not stated if ipsilateral, or bilateral, or contralateral AND not stated if single or multiple	*	D	D
60	Stated as N2NOS	N2NOS	RN	RN
70	Stated as N3, no other information	N3	RN	RN
80	Lymph nodes, NOS, no other information	*	RN	RN
99	Unknown; not stated Regional lymph node(s) cannot be assessed Not documented in patient record	NX	U	U

Subglottic Larynx
Reg LN Pos
SEE STANDARD TABLE

Subglottic Larynx
Reg LN Exam
SEE STANDARD TABLE

Subglottic Larynx
CS Mets at DX (Revised: 08/14/2006)
Note: Supraclavicular and transverse cervical lymph nodes are now coded in CS Lymph Nodes because they are categorized as N rather than M in AJCC TNM. Any cases coded to 10 or 50 can be reviewed and recoded. The volume of cases affected should be small.

Code	Description	TNM	SS77	SS2000
00	No; none	M0	NONE	NONE
10	Distant lymph node(s) Mediastinal Distant lymph node(s), NOS	M1	D	D
40	Distant metastases except distant lymph node(s) (code 10) Distant metastasis, NOS Carcinomatosis	M1	D	D
50	(10) + (40) Distant lymph node(s) plus other distant metastases	M1	D	D
99	Unknown if distant metastasis Distant metastasis cannot be assessed Not documented in patient record	MX	U	U

Collaborative Staging Codes
Larynx, Overlapping Lesion or Not Otherwise Specified
C32.3, C32.8-C32.9
C32.3 Laryngeal cartilage
C32.8 Overlapping lesion of larynx
C32.9 Larynx, NOS

Larynx, Overlapping Lesion or Not Otherwise Specified
CS Tumor Size
SEE STANDARD TABLE

Larynx, Overlapping Lesion or Not Otherwise Specified
CS Extension (Revised: 03/17/2004)

Code	Description	TNM	SS77	SS2000
00	In situ; noninvasive	Tis	IS	IS
10	Invasive tumor confined to site of origin	T1	L	L
20	Tumor involves more than one subsite, WITHOUT fixation or NOS	T2	L	L
30	Tumor involves adjacent regions(s) of larynx	T2	L	L
35	Impaired vocal cord mobility	T2	L	L
40	Tumor limited to larynx WITH vocal cord fixation	T3	L	L
45	Localized, NOS	T1	L	L
60	Hypopharynx, NOS Postcricoid area Pre-epiglottic tissues Pyriform sinus (pyriform fossa) Vallecula	T3	RE	RE
68	Extension to/through cricoid cartilage and thyroid cartilage	T4a	RE	D
70	Extension to/through: Cervical esophagus Deep muscle of tongue Extrinsic (strap) muscles Omohyoid Sternohyoid Sternothyroid Thyrohyoid	T4a	D	D

70 cont'd	Oropharynx Skin Soft tissues of neck Thyroid gland Trachea	T4a	D	D
80	Further contiguous extension, including: Mediastinal structures Prevertebral space Tumor encases carotid artery	T4b	D	D
95	No evidence of primary tumor	T0	U	U
99	Unknown extension Primary tumor cannot be assessed Not documented in patient record	TX	U	U

Larynx, Overlapping Lesion or Not Otherwise Specified
CS TS/Ext-Eval
SEE STANDARD TABLE
NOTE: *For these primary sites code CS Tumor Size/Ext Eval on the basis of the CS Extension field only.*

Larynx, Overlapping Lesion or Not Otherwise Specified
CS Lymph Nodes (Revised: 08/21/2006)
Note 1: For head and neck schemas, this field includes all lymph nodes defined as Levels I-VII and Other by AJCC.
Note 2: For head and neck schemas, additional information about lymph nodes (size of involved nodes, extracapsular extension, and levels involved) is coded in Site-Specific Factors 1-6.
Note 3: If laterality of lymph nodes is not specified, assume nodes are ipsilateral. Midline nodes are considered ipsilateral.
Note 4: For head and neck cancers, if lymph nodes are described only as "supraclavicular", try to determine if they are in Level IV (deep to the sternocleidomastoid muscle, in the lower jugular chain) or Level V (in the posterior triangle, inferior to the transverse cervical artery) and code appropriately. If the specific level cannot be determined, consider them as Level V nodes.

Code	Description	TNM	SS77	SS2000
00	None; no regional lymph node involvement	N0	NONE	NONE
10	Single positive ipsilateral regional node: Level II Jugulodigastric (subdigastric) Upper deep cervical Upper jugular Level III Middle deep cervical	*	RN	RN

10 cont'd	Mid-jugular Level IV Jugulo-omohyoid (supraomohyoid) Lower deep cervical Lower jugular Level VI Anterior deep cervical Delphian node Laterotracheal Paralaryngeal Paratracheal Prelaryngeal Pretracheal Recurrent laryngeal Cervical, NOS Deep cervical, NOS Internal jugular, NOS Regional lymph node, NOS Stated as N1, NOS	*	RN	RN
11	Single positive ipsilateral regional node: Level I Submandibular (submaxillary) Submental Other groups Retropharyngeal Mandibular, NOS	*	D	RN
12	Single positive ipsilateral regional node: Level V node Posterior cervical Posterior triangle (spinal accessory and transverse cervical) (upper, middle, and lower corresponding to the levels that define upper, middle, and lower jugular nodes) Level VII node Upper mediastinum (for other mediastinal nodes see CS Mets at DX) Other groups Intraparotid Parapharyngeal Periparotid Sub-occipital Supraclavicular, NOS (See Note 4)	*	D	RN

18	Stated as N1, no other information	N1	RN	RN
19	Stated as N2a, no other information	N2a	RN	RN
20	Multiple positive ipsilateral nodes listed in code 10	*	RN	RN
21	Multiple positive ipsilateral nodes listed in code 11	*	D	RN
22	Multiple positive ipsilateral nodes listed in code 12	*	D	D
29	Stated as N2b, no other information	N2b	RN	RN
30	Regional lymph nodes as listed in code 10: Positive ipsilateral node(s), not stated if single or multiple	*	RN	RN
31	Regional lymph nodes as listed in code 11: Positive ipsilateral node(s), not stated if single or multiple	*	D	RN
32	Regional lymph nodes as listed in code 12: Positive ipsilateral node(s), not stated if single or multiple	*	D	D
40	Regional lymph nodes as listed in code 10: Positive bilateral or contralateral nodes	*	RN	RN
41	Regional lymph nodes as listed in code 11: Positive bilateral or contralateral nodes	*	D	RN
42	Regional lymph nodes as listed in code 12: Positive bilateral or contralateral nodes	*	D	D
49	Stated as N2c, no other information	N2c	RN	RN
50	Regional lymph nodes as listed in code 10: Positive node(s) not stated if ipsilateral, or bilateral, or contralateral AND not stated if single or multiple	*	RN	RN
51	Regional lymph nodes as listed in code 11: Positive node(s) not stated if ipsilateral, or bilateral, or contralateral AND not stated if single or multiple	*	D	RN
52	Regional lymph nodes as listed in code 12: Positive node(s), not stated if ipsilateral, or bilateral, or contralateral AND not stated if single or multiple	*	D	D
60	Stated as N2, NOS	N2NOS	RN	RN
70	Stated as N3, no other information	N3	RN	RN

80	Lymph nodes, NOS, no other information	*	RN	RN
99	Unknown; not stated Regional lymph node(s) cannot be assessed Not documented in patient record	NX	U	U

Larynx, Overlapping Lesion or Not Otherwise Specified
Reg LN Pos
SEE STANDARD TABLE

Larynx, Overlapping Lesion or Not Otherwise Specified
Reg LN Exam
SEE STANDARD TABLE

Larynx, Overlapping Lesion or Not Otherwise Specified
CS Mets at DX (Revised: 08/14/2006)
Note: Supraclavicular and transverse cervical lymph nodes are now coded in CS Lymph Nodes because they are categorized as N rather than M in AJCC TNM. Any cases coded to 10 or 50 can be reviewed and recoded. The volume of cases affected should be small.

Code	Description	TNM	SS77	SS2000
00	No; none	M0	NONE	NONE
10	Distant lymph node(s) Mediastinal Distant lymph node(s), NOS	M1	D	D
40	Distant metastases except distant lymph node(s) (code 10) Distant metastasis, NOS Carcinomatosis	M1	D	D
50	(10) + (40) Distant lymph node(s) plus other distant metastases	M1	D	D
99	Unknown if distant metastasis Distant metastasis cannot be assessed Not documented in patient record	MX	U	U

Site Specific Surgery Codes
Larynx
C320–C329
(Except for M9750, 9760-9764, 9800-9820, 9826, 9831-9920, 9931-9964, 9980-9989)

Codes

00 None; no surgery of primary site; autopsy ONLY

10 Local tumor destruction, NOS
- 11 Photodynamic therapy (PDT)
- 12 Electrocautery; fulguration (includes use of hot forceps for tumor destruction)
- 13 Cryosurgery
- 14 Laser
- 15 Stripping

No specimen sent to pathology from surgical events 10–15

20 Local tumor excision, NOS
- 26 Polypectomy
- 27 Excisional biopsy

Any combination of 20 or 26–27 WITH
- 21 Photodynamic therapy (PDT)
- 22 Electrocautery
- 23 Cryosurgery
- 24 Laser ablation

[**SEER Note**: Codes 21 to 24 above combine 20 Local tumor excision, 26 Polypectomy or 27 Excisional biopsy with 21 PDT, 22 Electrocautery, 23 Cryosurgery, or 24 Laser ablation]
- 25 Laser excision
- 28 Stripping

Specimen sent to pathology from surgical events 20–28

30 Partial excision of the primary site, NOS; subtotal/partial laryngectomy NOS; hemilaryngectomy NOS
- 31 Vertical laryngectomy
- 32 Anterior commissure laryngectomy
- 33 Supraglottic laryngectomy

[**SEER Notes**: Vertical laryngectomy: Removal of involved true vocal cord, ipsilateral false vocal cord, intervening ventricle, ipsilateral thyroid and may include removal of the arytenoids. Supraglottic laryngectomy: Conservative surgery intended to preserve the laryngeal function. Standard procedure involves removal of epiglottis, false vocal cords, aryepiglottic folds, arytenoid cartilages, ventricle, upper one third of thyroid cartilage, thyroid membrane. The true vocal cords and arytenoids remain in place to allow vocalization and deglutition.]

40 Total or radical laryngectomy, NOS
- 41 Total laryngectomy ONLY
- 42 Radical laryngectomy ONLY

[**SEER Note**: Radical laryngectomy: Includes removal of adjacent sites. Do not code the removal of adjacent sites in Surgical Procedure of Other Site.]

50 Pharyngolaryngectomy

80 Laryngectomy, NOS

90 Surgery, NOS

99 Unknown if surgery performed; death certificate ONLY