

Neuroendocrine Tumors of Colon**C18.0, C18.2-C18.9 (excluding Appendix)**

(M-8153, 8240-8242, 8246, 8249)

C18.0 Cecum

C18.2 Ascending colon

C18.3 Hepatic flexure of colon

C18.4 Transverse colon

C18.5 Splenic flexure of colon

C18.6 Descending colon

C18.7 Sigmoid colon

C18.8 Overlapping lesion of colon

C18.9 Colon, NOS

Note 1: For this schema, AJCC only stages well-differentiated neuroendocrine tumors. Please note that the "concept" of well-differentiated is reflected in the histology code. The grade code is not needed in order to select the correct schema, but does need to be coded.

Note 2: This schema is also used for carcinoid tumors and malignant gastrinomas.

NETColon**CS Tumor Size**

Note: Code the specific tumor size as documented in the medical record. If the ONLY information regarding tumor size is the physician's statement of the "T" category, assign code 011 (T1, NOS), 021 (T2, NOS) or 999 (T3, NOS or T4, NOS) . (Refer to the CS Extension table for instructions on coding extension).

Code	Description
000	No mass/tumor found
001-988	001 - 988 millimeters (code exact size in millimeters)
989	989 millimeters or larger
990	Microscopic focus or foci only, no size of focus given
991	Described as "less than 1 cm"
992	OBSOLETE DATA RETAINED V0200 Described as "less than 2 cm," or "greater than 1 cm," or "between 1 cm and 2 cm"
993	OBSOLETE DATA RETAINED V0200 Described as "less than 3 cm," or "greater than 2 cm," or "between 2 cm and 3 cm"

Code	Description
994	OBSOLETE DATA RETAINED V0200 Described as "less than 4 cm," or "greater than 3 cm," or "between 3 cm and 4 cm"
995	OBSOLETE DATA RETAINED V0200 Described as "less than 5 cm," or "greater than 4 cm," or "between 4 cm and 5 cm"
998	OBSOLETE DATA RETAINED V0200 Familial/multiple polyposis (M-8220/8221)
999	Unknown; size not stated Not documented in patient record

NETColon CS Extension

Note 1: Ignore intraluminal extension to adjacent segment(s) of colon/rectum or to the ileum from the cecum; code depth of invasion or extracolonic spread as indicated.

Note 2: Codes 600-800 are used for contiguous extension from the site of origin. Discontinuous involvement is coded in CS Mets at DX.

Code	Description	TNM 7 Map	TNM 6 Map	SS77 Map	SS2000 Map
000	In situ; noninvasive; intraepithelial	^	Tis	IS	IS
050	(Adeno)carcinoma in a polyp or adenoma, noninvasive	^	Tis	IS	IS
100	Invasive tumor confined to mucosa, NOS (including intramucosal, NOS)	^	Tis	L	L
110	Lamina propria, including lamina propria in the stalk of a polyp	^	Tis	L	L
120	Confined to and not through the muscularis mucosae, including muscularis mucosae in the stalk of a polyp.	^	Tis	L	L
130	Confined to head of polyp, NOS	^	T1	L	L

Code	Description	TNM 7 Map	TNM 6 Map	SS77 Map	SS2000 Map
140	Confined to stalk of polyp, NOS	^	T1	L	L
150	Invasive tumor in polyp, NOS	^	T1	L	L
160	Invades submucosa (superficial invasion), including submucosa in the stalk of a polyp	^	T1	L	L
170	Stated as T1a with no other information on extension	^	T1	L	L
180	Stated as T1b with no other information on extension	^	T1	L	L
190	Stated as T1[NOS] with no other information on extension	^	T1	L	L
200	Muscularis propria invaded	^	T2	L	L
210	Stated as T2, NOS with no other information on extension	^	T2	L	L
300	Localized, NOS Confined to colon, NOS	^	T1	L	L
400	Extension through wall, NOS Invasion through muscularis propria or muscularis, NOS Non-peritonealized pericolic tissues invaded Perimuscular tissue invaded Subserosal tissue/(sub)serosal fat invaded Transmural, NOS	T3	T3	L	L
410	Stated as T3, NOS with no other information on extension	T3	T3	L	L
420	Fat, NOS	T3	T3	RE	RE

Code	Description	TNM 7 Map	TNM 6 Map	SS77 Map	SS2000 Map
450	Extension to: All colon sites: Adjacent tissue(s), NOS Connective tissue Mesenteric fat Mesentery Mesocolon Pericolonic fat Ascending and descending colon Retroperitoneal fat Transverse colon/flexures Gastrocolic ligament Greater omentum	T3	T3	RE	RE
460	Adherent to other organs or structures, but no microscopic tumor found in adhesion(s)	T3	T3	RE	RE
490	Stated as T4, NOS with no other information on extension	T4	T4	RE	RE
500	Invasion of/through serosa (mesothelium) (visceral peritoneum)	T4	T4	RE	RE
550	Any of [(420) to (450)] + (500)	T4	T4	RE	RE
570	Adherent to other organs or structures, NOS	T4	T4	RE	RE
600	All colon sites: Small intestine Cecum: Greater omentum Ascending colon: Greater omentum Liver, right lobe Transverse colon and flexures: Gallbladder/bile ducts Kidney Liver Pancreas	T4	T4	RE	RE

Code	Description	TNM 7 Map	TNM 6 Map	SS77 Map	SS2000 Map
600 cont'd	Spleen Stomach Descending colon: Greater omentum Pelvic wall Spleen Sigmoid colon: Greater omentum Pelvic wall	T4	T4	RE	RE
650	All colon sites: Abdominal wall Retroperitoneum (excluding fat)	T4	T4	RE	RE
660	Ascending colon: Right kidney Right ureter Descending colon: Left kidney Left ureter	T4	T4	RE	RE
700	Cecum, ascending, descending and sigmoid colon: Fallopian tube Ovary Uterus	T4	T4	D	D
750	All colon sites unless otherwise stated above: Adrenal (suprarenal) gland Bladder Diaphragm Fistula to skin Gallbladder Other segment(s) of colon via serosa	T4	T4	D	D
800	Further contiguous extension: Cecum: Kidney Liver Ureter	T4	T4	D	D

Code	Description	TNM 7 Map	TNM 6 Map	SS77 Map	SS2000 Map
800 cont'd	Transverse colon and flexures: Ureter Sigmoid colon: Cul de sac (rectouterine pouch) Ureter Other contiguous extension	T4	T4	D	D
950	No evidence of primary tumor	T0	T0	U	U
999	Unknown extension Primary tumor cannot be assessed Not documented in patient record	TX	TX	U	U

^ For codes 000-300 ONLY, the T category is assigned based on the value of CS Tumor Size, as shown in the Extension Size 7th Table for this site.

NETColon

CS Tumor Size/Ext Eval

See Standard Table

NETColon

CS Lymph Nodes

Note 1: Code only regional nodes and nodes, NOS, in this field. Distant nodes are coded in the field Mets at DX.

Note 2: Inferior mesenteric nodes are coded in CS Mets at DX for cecum, ascending colon, transverse colon, and hepatic flexure. Superior mesenteric nodes are coded in CS Mets at DX for all colon sites.

Code	Description	TNM 7 Map	TNM 6 Map	SS77 Map	SS2000 Map
000	None; no regional lymph node involvement	N0	N0	NONE	NONE
100	Regional lymph nodes for all colon sites: Colic (NOS) Epicolic (adjacent to bowel wall) Mesocolic (NOS) Paracolic/pericolic	N1	*	RN	RN
200	Regional lymph nodes, for specific	N1	*	RN	RN

Code	Description	TNM 7 Map	TNM 6 Map	SS77 Map	SS2000 Map
200 cont'd	subsites: Cecum: Cecal: anterior (prececal), posterior (retrocecal); NOS Ileocolic Right colic Ascending colon: Ileocolic Middle colic Right colic Transverse colon and flexures: Inferior mesenteric for splenic flexure only Left colic for splenic flexure only Middle colic Right colic for hepatic flexure only Descending colon: Inferior mesenteric Left colic Sigmoid Sigmoid colon: Inferior mesenteric Sigmoidal (sigmoid mesenteric) Superior hemorrhoidal Superior rectal	N1	*	RN	RN
300	Regional lymph nodes for all colon sites: Mesenteric, NOS Regional lymph node(s), NOS	N1	*	RN	RN
400	Stated as N1, NOS	N1	N1	RN	RN
450	OBSOLETE DATA RETAINED V0200 Stated as N2 pathologic	ERROR	N2	RN	RN
800	Lymph nodes, NOS	N1	*	RN	RN

Code	Description	TNM 7 Map	TNM 6 Map	SS77 Map	SS2000 Map
999	Unknown; not stated Regional lymph node(s) cannot be assessed Not documented in patient record	NX	NX	U	U

* For codes 100-300 and 800 ONLY: when CS Lymph Nodes Eval is 0, 1, 5, or 9, the N category is assigned from the Lymph Nodes Clinical Evaluation 6th Table, using Reg LN Pos and CS Site-Specific Factor 2; when CS Regional Nodes Eval is 2, 3, 6, 8, or not coded, the N category is determined from the Lymph Nodes Pathologic Evaluation 6th Table Also Used When CS Reg Nodes Eval is Not Coded using Reg LN Pos.

NETColon**Reg LN Pos**

Note: Record this field even if there has been preoperative treatment.

See Standard Table

NETColon**Reg LN Exam**

See Standard Table

NETColon**CS Mets at DX**

Code	Description	TNM 7 Map	TNM 6 Map	SS77 Map	SS2000 Map
00	No; none	M0	M0	NONE	NONE
08	Cecum, ascending, hepatic flexure and transverse colon: Superior mesenteric lymph nodes only	M1	M1	RN	D
10	Distant lymph node(s) other than code 08 For all colon sites: Common iliac Distant lymph node(s), NOS External iliac	M1	M1	D	D

Code	Description	TNM 7 Map	TNM 6 Map	SS77 Map	SS2000 Map
10 cont'd	Para-aortic Retroperitoneal For cecum, ascending colon, transverse colon, and hepatic flexure: Inferior mesenteric For splenic flexure, descending colon, and sigmoid colon: Superior mesenteric	M1	M1	D	D
40	Distant metastases except distant lymph node(s) (codes 08-10) Carcinomatosis	M1	M1	D	D
50	(40) + ((08) or (10)) Distant lymph node(s) plus other distant metastases	M1	M1	D	D
60	Distant metastasis, NOS M1, NOS	M1	M1	D	D
99	Unknown if distant metastasis Distant metastasis cannot be assessed Not documented in patient record	M0	MX	U	U