

Ampulla of Vater**C24.1**

C24.1 Ampulla of Vater

Ampulla Vater**CS Tumor Size**

See Standard Table

Ampulla Vater**CS Extension**

Code	Description	TNM 7 Map	TNM 6 Map	SS77 Map	SS2000 Map
000	In situ; non-invasive; intraepithelial Stated as Tis, NOS	Tis	Tis	IS	IS
100	Invasive tumor confined/limited to ampulla of Vater or extending to sphincter of Oddi	T1	T1	L	L
300	Localized, NOS Stated as T1, NOS	T1	T1	L	L
420	Duodenal wall Stated as T2, NOS	T2	T2	RE	RE
520	Pancreas Stated as T3, NOS	T3	T3	RE	RE
620	Common bile duct	T4	T4	RE	RE
650	Extrahepatic bile ducts other than common bile duct or sphincter of Oddi	T4	T4	RE	RE
700	Extension to other adjacent organs or tissues: Blood vessels(major): Hepatic artery Portal vein Gallbladder Hepatic flexure Lesser omentum Liver including porta hepatis Peripancreatic soft tissues	T4	T4	RE	RE

Code	Description	TNM 7 Map	TNM 6 Map	SS77 Map	SS2000 Map
700 cont'd	Stomach, NOS: Distal Transverse colon	T4	T4	RE	RE
750	Stomach, proximal	T4	T4	RE	D
780	Stated as T4, NOS	T4	T4	D	D
800	Further contiguous extension Other adjacent organs	T4	T4	D	D
950	No evidence of primary tumor	T0	T0	U	U
999	Unknown extension Primary tumor cannot be assessed Not documented in patient record	TX	TX	U	U

Ampulla Vater**CS Tumor Size/Ext Eval**

See Standard Table

Ampulla Vater**CS Lymph Nodes**

Note 1: Code only regional nodes and nodes, NOS, in this field. Distant nodes are coded in the field Mets at DX.

Note 2: Splenic lymph nodes and those located at the tail of the pancreas are not considered regional and should be coded under Mets at DX.

Code	Description	TNM 7 Map	TNM 6 Map	SS77 Map	SS2000 Map
000	No regional lymph node involvement Stated as N0, NOS	N0	N0	NONE	NONE
100	Regional lymph nodes: Hepatic Hepatic artery Node of the foramen of Winslow (epiploic) (omental) Pancreaticoduodenal	N1	N1	RN	RN

Code	Description	TNM 7 Map	TNM 6 Map	SS77 Map	SS2000 Map
100 cont'd	Peripancreatic (except at tail of pancreas, see CS Mets at DX) Periportal Lymph nodes: Anterior to the ampulla of Vater Inferior to the ampulla of Vater Posterior to the ampulla of Vater Superior to the ampulla of Vater Regional lymph node(s), NOS Stated as N1, NOS	N1	N1	RN	RN
110	Regional lymph nodes: Celiac Infrapyloric (subpyloric) Lateral aortic (lumbar) Proximal mesenteric Retroperitoneal Superior mesenteric	N1	N1	D	RN
800	Lymph nodes, NOS	N1	N1	RN	RN
999	Unknown; not stated Regional lymph nodes cannot be assessed Not documented in patient record	NX	NX	U	U

Ampulla Vater**Reg LN Pos**

Note: Record this field even if there has been preoperative treatment.

See Standard Table

Ampulla Vater**Reg LN Exam**

See Standard Table

**Ampulla Vater
CS Mets at DX**

Code	Description	TNM 7 Map	TNM 6 Map	SS77 Map	SS2000 Map
00	No; none	M0	M0	NONE	NONE
10	Distant lymph node(s) Node(s) at the tail of the pancreas Para-aortic Splenic lymph node(s) Distant lymph nodes, NOS	M1	M1	D	D
40	Distant metastases except distant lymph node(s) (code 10) Carcinomatosis (Includes seeding of peritoneum, even if limited to the lesser sac region; positive peritoneal cytology)	M1	M1	D	D
50	(10) + (40) Distant lymph node(s) plus other distant metastases	M1	M1	D	D
60	Distant metastasis, NOS Stated as M1, NOS	M1	M1	D	D
99	Unknown if distant metastasis Distant metastasis cannot be assessed Not documented in patient record	M0	MX	U	U