

Corpus Carcinoma**Carcinoma and Carcinosarcoma of Corpus Uteri; Uterus, NOS (excluding Placenta and Adenosarcoma, Leiomyosarcoma, and Endometrial Stromal Sarcoma (ESS))****C54.0-C54.3, C54.8-C54.9, C55.9****M-8000-8790, 8980-8981, 9700-9701****C54.0 Isthmus uteri****C54.1 Endometrium****C54.2 Myometrium****C54.3 Fundus uteri****C54.8 Overlapping lesion of corpus uteri****C54.9 Corpus uteri****C55.9 Uterus, NOS**

Note 1: AJCC 7th Edition TNM staging reflects the new staging adopted by the International Federation of Gynecology and Obstetrics (FIGO) and utilizes three new staging schemas for cancer of the Corpus Uteri based on histology. This is a change from the AJCC 6th Edition TNM staging. The three new schemas are: 1. Carcinoma and carcinosarcoma 2. Leiomyosarcoma and endometrial stromal sarcoma (ESS) 3. Adenosarcoma

Note 2: Carcinosarcoma should be staged as carcinoma.

Note 3: This schema is for carcinoma and carcinosarcoma. When coding pay attention to the FIGO Editions and descriptions about cell and tumor types, disease extension, lymph node status and metastasis.

Note 4: AJCC TNM values correspond to the stages accepted by the Federation Internationale de Gynecologic et d'Obstetrique (FIGO). Note that FIGO uses a single stage value which corresponds to different groupings of T, N, and M values.

Corpus Carcinoma**CS Tumor Size****See Standard Table****Corpus Carcinoma****CS Extension**

Note 1. When both FIGO stage and extension detail are available, record the code with extension detail in preference to a statement of FIGO stage.

Note 2: FIGO no longer includes Stage 0 (Tis) for this site.

Note 3: To assign a T2 category, stromal connective tissue of the cervix must be involved; endocervical glandular involvement only is not considered a T2 tumor in AJCC 7. Use 123, 133, 143, or 145 if endometrial or myometrial involvement is described.

Note 4: FIGO IIIA and IIIB are extension, whereas FIGO IIIC is nodal involvement. FIGO IVA is extension, whereas FIGO IVB is distant metastasis. FIGO Stage III, NOS and FIGO Stage IV, NOS are assigned based on tumor extension and/or metastasis. Code CS Extension based on a physician's statement of FIGO Stage III, NOS or FIGO Stage IV, NOS only if it is known that the stage value is assigned because of tumor extension.

Note 5: Positive cytology is reported separately (CS Site Specific Factor 2), and is not included as a staging element for AJCC 7 staging. It is included as a staging element for AJCC 6 and Summary Stage. Since "cancer cells in ascites or in peritoneal washings" was not specifically categorized in the

1977 Summary Staging Guide, it is unclear to which stage previous cases may have been coded in that staging system.

Note 6: According to the AJCC, extension to the bowel or bladder mucosa must be proven by biopsy in order to rule out bullous edema.

| Code | Description | TNM 7 Map | TNM 6 Map | SS77 Map | SS2000 Map |
|------|---|-----------|-----------|----------|------------|
| 000 | In situ, intraepithelial, noninvasive, preinvasive | Tis | Tis | IS | IS |
| 100 | Invasive cancer confined to corpus uteri | T1NOS | * | ** | ** |
| 110 | Confined to endometrium (stroma) | T1a | * | ** | ** |
| 120 | Tumor invades less than one-half of myometrium Invasion of inner half of myometrium | T1a | * | ** | ** |
| 123 | Endocervical glandular involvement WITH tumor limited to endometrium or invading less than one-half of myometrium (See Note 3) | T1a | * | RE | RE |
| 125 | FIGO Stage IA | T1a | * | ** | ** |
| 130 | Tumor invades one-half or more of myometrium Invasion of outer half of myometrium (See Note 4) | T1b | * | ** | ** |
| 133 | Endocervical glandular involvement WITH tumor invading one-half or more of myometrium (See Note 3) | T1b | * | RE | RE |
| 135 | FIGO Stage IB | T1b | * | ** | ** |
| 140 | Invasion of myometrium, NOS | T1NOS | * | ** | ** |
| 143 | Endocervical glandular involvement WITH tumor invading myometrium, NOS | T1NOS | * | RE | RE |

| Code | Description | TNM 7 Map | TNM 6 Map | SS77 Map | SS2000 Map |
|------|--|-----------|-----------|----------|------------|
| 145 | Endocervical glandular involvement and involvement of myometrium not specified | T1NOS | * | RE | RE |
| 160 | OBsolete DATA RETAINED AND REVIEWED V0203 See codes 540 and 545 Tunica serosa of the visceral peritoneum (serosa covering the corpus) | T1NOS | T3a | ** | ** |
| 180 | FIGO Stage I [NOS] | T1NOS | * | ** | ** |
| 400 | Localized, NOS | T1NOS | * | ** | ** |
| 500 | Cervix uteri, NOS, but not beyond uterus | T2 | * | RE | RE |
| 510 | OBsolete DATA RETAINED AND REVIEWED V0203 See code 123, 133, 143, and 145 Endocervical glandular involvement only | T1NOS | * | RE | RE |
| 520 | Cervical stromal invasion | T2 | T2b | RE | RE |
| 525 | FIGO Stage II | T2 | * | RE | RE |
| 540 | Extension or metastasis to tunica serosa (visceral peritoneum of corpus, serosa covering the corpus) WITHOUT involvement of cervix | T3a | T3a | ** | ** |
| 545 | Extension or metastasis to tunica serosa of corpus WITH involvement of cervix | T3a | T3a | RE | RE |
| 550 | Extension or metastasis to: Adnexa: Fallopian tube(s), Ovary(ies) | T3a | T3a | RE | RE |

| Code | Description | TNM 7 Map | TNM 6 Map | SS77 Map | SS2000 Map |
|------|--|-----------|-----------|----------|------------|
| 600 | <p>OBSELETE DATA RETAINED AND REVIEWED V0203 See codes 550, 655, and 800 Extension or metastasis within true pelvis:</p> <ul style="list-style-type: none"> Adnexa Fallopian tube(s) Ligaments: Broad, round, uterosacral Ovary(ies) Parametrium Pelvic serosa Tunica serosa (parietal lining of the pelvic or abdominal cavity) | T3a | T3a | RE | RE |
| 610 | <p>OBSELETE DATA RETAINED V0200; See Note 5 and CS Site-Specific Factor 2 Cancer cells in ascites Cancer cells in peritoneal washings FIGO Stage IIIA</p> | ERROR | T3a | RE | RE |
| 620 | <p>OBSELETE DATA CONVERTED V0200; (See code 645) Ureter and vulva</p> | ERROR | ERROR | ERROR | ERROR |
| 630 | FIGO Stage IIIA | T3a | T3a | RE | RE |
| 635 | <p>Extension or metastasis to:</p> <ul style="list-style-type: none"> Ligaments: Broad Round Parametrium, NOS Visceral peritoneum of pelvic organs excluding serosa of corpus | T3b | T3b | RE | RE |
| 640 | Extension or metastasis to vagina | T3b | T3b | D | RE |
| 645 | <p>Extension to:</p> <ul style="list-style-type: none"> Ureter Vulva | T3b | T3b | D | RE |

| Code | Description | TNM 7 Map | TNM 6 Map | SS77 Map | SS2000 Map |
|------|--|-----------|-----------|----------|------------|
| 650 | OBSOLETE DATA RETAINED AND REVIEWED V0203 See codes 655 and 662 Extension or metastasis to pelvic wall(s) Described clinically as "frozen pelvis", NOS | T3b | T3b | RE | RE |
| 655 | Extension or metastasis to: Pelvic wall(s) Parietal serosa of pelvic wall | T3b | T3b | RE | RE |
| 660 | Extension or metastasis to: Bladder wall Bladder, NOS excluding mucosa Rectal wall Rectum, NOS excluding mucosa | T3b | T3b | RE | RE |
| 662 | Described clinically as "frozen pelvis", NOS | T3b | T3b | D | RE |
| 663 | (660 or 662) + (640 or 645) | T3b | T3b | D | RE |
| 665 | FIGO Stage IIIB | T3b | T3b | RE | RE |
| 670 | OBSOLETE DATA CONVERTED V0203 ; See code 663 (662 or 660) + (645 or 640) | ERROR | ERROR | ERROR | ERROR |
| 680 | FIGO Stage III [NOS] based on tumor extension (See Note 4.) | T3NOS | T3NOS | RE | RE |
| 700 | OBSOLETE DATA CONVERTED V0200 ; (See code 710) Extension to bowel mucosa or bladder mucosa (excluding bullous edema) FIGO Stage IVA FIGO Stage IV, NOS | ERROR | ERROR | ERROR | ERROR |

| Code | Description | TNM 7 Map | TNM 6 Map | SS77 Map | SS2000 Map |
|------|---|-----------|-----------|----------|------------|
| 710 | OBsolete DATA REVIEWED AND CHANGED V0203 See codes 715, 810, and 820 Extension to bowel mucosa or bladder mucosa (excluding bullous edema) Note: On conversion of code 700 to 710, code 710 also included FIGO Stage IVA, FIGO Stage IV, NOS | ERROR | ERROR | ERROR | ERROR |
| 715 | Extension to bowel mucosa or bladder mucosa (excluding bullous edema) | T4 | T4 | D | D |
| 800 | Further contiguous extension Abdominal serosa (visceral or parietal peritoneum of abdomen) Cul de sac (rectouterine pouch or Pouch of Douglas) Sigmoid colon Small intestine | T4 | T4 | D | D |
| 810 | FIGO Stage IVA | T4 | T4 | D | D |
| 820 | FIGO Stage IV [NOS] based on tumor extension (See Note 4.) | T4 | T4 | D | D |
| 950 | No evidence of primary tumor | T0 | T0 | U | U |
| 999 | Unknown; extension not stated Primary tumor cannot be assessed Not documented in patient record | TX | TX | U | U |

* For CS Extension codes 100-145, 180-510, and 525 ONLY. If it is a CS Version 1 case, (CS Version Input Original equals 01XXXX and Year of Diagnosis is before 2010), the T category for AJCC 6 stages is assigned based on the value of CS Extension and peritoneal cytology coded in CS Site-Specific Factor 2 as shown in the Extension Cytology AJCC 6 Table CSv1. If it is a CS Version 2 case (CS Version Input Original equals 02XXXX or Year of Diagnosis is greater than 2009), the T category for AJCC 6 staging is based on the value of CS Extension and peritoneal cytology coded in CS Site-Specific Factor 2 as shown in the Extension Cytology AJCC 6 Table CSv2.

** For CS Extension codes 100-120, 125-130, 135-140, 160-400, and 540 ONLY. The SS77 and SS2000 stages are assigned based on the value of CS Extension and peritoneal cytology status coded in CS Site-Specific Factor 2 as shown in the Extension Cytology Summary Stage Table.

Corpus Carcinoma
CS Tumor Size/Ext Eval
See Standard Table

Corpus Carcinoma
CS Lymph Nodes

Note 1: Code only regional nodes and nodes, NOS, in this field. Distant nodes are coded in CS Mets at DX. Regional nodes include bilateral and contralateral involvement of named nodes.

Note 2: When both FIGO stage and regional lymph node detail are available, record the code with regional lymph node detail in preference to a statement of FIGO stage.

Note 3: If either exploratory or definitive surgery is done with no mention of lymph nodes, assume nodes are negative.

Note 4: If the clinician says "adnexa palpated" but does not mention lymph nodes, assume lymph nodes are not involved.

| Code | Description | TNM 7 Map | TNM 6 Map | SS77 Map | SS2000 Map |
|------|---|-----------|-----------|----------|------------|
| 000 | No regional lymph node involvement | N0 | N0 | NONE | NONE |
| 100 | Regional lymph node(s): Iliac, NOS: Common External Internal (hypogastric) Obturator Paracervical Parametrial Sacral, NOS: Lateral (laterosacral) Middle (promontorial) (Gerota's node) Presacral Uterosacral | N1 | N1 | RN | RN |
| 140 | Pelvic lymph node(s), NOS | N1 | N1 | RN | RN |
| 160 | FIGO Stage IIIC1 | N1 | N1 | RN | RN |
| 200 | Regional lymph node(s): Aortic, NOS: Lateral (lumbar) Para-aortic Periaortic | N2 | N1 | RN | RN |

| Code | Description | TNM 7 Map | TNM 6 Map | SS77 Map | SS2000 Map |
|------|---|-----------|-----------|----------|------------|
| 210 | 200 + (140 or 100) | N2 | N1 | RN | RN |
| 250 | FIGO Stage IIIC2 | N2 | N1 | RN | RN |
| 500 | FIGO Stage IIIC [NOS] | N1 | N1 | RN | RN |
| 800 | Regional lymph node(s), NOS | N1 | N1 | RN | RN |
| 999 | Unknown; regional lymph nodes not stated; Regional lymph node(s) cannot be assessed; Not documented in patient record | NX | NX | U | U |

**Corpus Carcinoma
CS Lymph Nodes Eval
See Standard Table**

**Corpus Carcinoma
Regional Nodes Positive
See Standard Table**

Note: Record this field even if there has been preoperative treatment.

**Corpus Carcinoma
Regional Nodes Examined
See Standard Table**

**Corpus Carcinoma
CS Mets at DX**

Note 1: Metastasis to adnexa, parametria, pelvic serosa, or vagina is coded in CS Extension.

Note 2: When both Federation Internationale de Gynecologie et d'Obstetrique (FIGO) stage and metastatic detail are available, record the code with metastatic detail in preference to a statement of FIGO stage.

Note 3: FIGO Stage IV, NOS is assigned based on tumor extension and/or metastasis. Code CS Mets at DX based on a physician's statement of FIGO Stage IV, NOS only if it is known that the stage value is assigned because of metastasis.

| Code | Description | TNM 7 Map | TNM 6 Map | SS77 Map | SS2000 Map |
|------|-----------------------|-----------|-----------|----------|------------|
| 00 | No distant metastasis | M0 | M0 | NONE | NONE |

| Code | Description | TNM 7 Map | TNM 6 Map | SS77 Map | SS2000 Map |
|------|--|-----------|-----------|----------|------------|
| 11 | Distant lymph node(s): Superficial inguinal | M1 | M1 | RN | D |
| 12 | Distant lymph node(s) other than code 11: Deep inguinal, NOS: Node of Cloquet or Rosenmuller (highest deep inguinal) Distant lymph node(s), NOS (Excluding metastasis to pelvic or para-aortic lymph nodes) | M1 | M1 | D | D |
| 40 | Distant metastases, except distant lymph node(s) (Excluding metastasis to vagina, pelvic, serosa, or adnexa) Carcinomatosis | M1 | M1 | D | D |
| 50 | 40 + (11 and/or 12) Distant metastasis plus distant lymph node(s) | M1 | M1 | D | D |
| 55 | FIGO Stage IVB | M1 | M1 | D | D |
| 60 | Distant metastasis, NOS | M1 | M1 | D | D |
| 70 | FIGO Stage IV [NOS] based on distant metastasis; (See note 3) | M1 | M1 | D | D |
| 99 | Unknown; distant metastasis not stated Distant metastasis cannot be assessed Not documented in patient record | M0 | MX | U | U |

Corpus Carcinoma**CS Mets Eval****See Standard Table**

Note: This item reflects the validity of the classification of the item CS Mets at DX only according to the diagnostic methods employed.

Corpus Carcinoma**CS Site-Specific Factor 2****Peritoneal Cytology****Note: See page A-132**

Note: Cytologic examination for malignant cells may be performed on ascites (fluid that has accumulated in the peritoneal cavity in excess amount) or the fluid (saline) that is introduced into the peritoneal cavity, and then removed by suction. The introduction of fluid may be termed peritoneal washing or peritoneal lavage.

| Code | Description |
|------|--|
| 000 | Negative |
| 010 | Positive, malignant cells positive |
| 020 | Test done, results suspicious or undetermined if negative or positive |
| 888 | OBSOLETE DATA CONVERTED V0200 See code 988 Not applicable for this site |
| 988 | Not applicable: Information not collected for this case; (May include cases converted from code 888 used in CSv1 for "Not applicable" or when the item was not collected. If this item is required to derive T, N, M, or any stage, use of code 988 may result in an error.) |
| 997 | Test ordered, results not in patient record |
| 998 | Test not done (test not ordered and not performed) No pathologic specimen available |
| 999 | Unknown or no information Not documented in patient record |

CorpusCarcinoma**CS Site-Specific Factor 3****Number of Positive Pelvic Nodes**

Note: Record the number of positive pelvic lymph nodes documented in the patient record. Apply the same instructions in General Rules, Part I for coding Regional Nodes Positive to this data item.

Note: See page A-67

| Code | Description |
|---------|--|
| 000 | All pelvic nodes examined negative. |
| 001-089 | 1 - 89 pelvic nodes positive (Exact number of nodes positive) |
| 090 | 90 or more pelvic nodes positive |
| 095 | Positive aspiration or core biopsy of pelvic lymph node(s) |
| 097 | Positive pelvic nodes, number not specified |
| 098 | No pelvic nodes examined |
| 888 | OBSOLETE DATA CONVERTED V0200 See code 988 Not applicable for this site |
| 988 | Not applicable: Information not collected for this case (May include cases converted from code 888 used in CSv1 for "Not applicable" or when the item was not collected. If this item is required to derive T, N, M, or any stage, use of code 988 may result in an error.) |
| 999 | Unknown if pelvic nodes are positive Not documented in patient record |